

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number  
**WO 01/96368 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 7/00**

(21) International Application Number: PCT/US01/19168

(22) International Filing Date: 14 June 2001 (14.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/211,892 14 June 2000 (14.06.2000) US  
60/213,387 23 June 2000 (23.06.2000) US

(71) Applicant (for all designated States except US): **CYTO-VAX BIOTECHNOLOGIES, INC.** [CA/CA]; Suite 308, 8925 - 51 Avenue, Edmonton, Alberta T6E 5J3 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HOUSTON, Michael, E.** [CA/CA]; 8767 Strathearn Drive, Edmonton, Alberta T6G 4C8 (CA). **HODGES, Robert, S.** [CA/US]; 6881 East - 12th Avenue, Denver, CA 80220 (US).

(74) Agents: **SWISS, Gerald, F.** et al.; Burns, Doane, Swecker & Mathis, LLP., P.O. Box 1404, Alexandria, VA 22313-1404 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF COILED-COIL STRUCTURAL SCAFFOLD TO GENERATE STRUCTURE-SPECIFIC PEPTIDES



(57) Abstract: This invention relates to the use of a coiled-coil structural scaffold to generate structure-specific peptides, including synthetic peptides derived from naturally occurring proteins of various origin. The structure of the synthetic peptides utilizes a scaffold of heptad repeat units into which epitopes from coiled-coil regions of native proteins are spliced. In particular, the synthetic peptides may be based on microbial proteins, especially surface proteins, which occur naturally in the coiled-coil form such as pneumococcal surface proteins A and C. The synthetic peptides are immunogenic and can be used to elicit an immune response in an animal. Accordingly, they are useful as vaccines or to stimulate antibody production or cell-mediated immunity to the naturally occurring protein.



WO 01/96368 A2

-1-

## USE OF COILED-COIL STRUCTURAL SCAFFOLD TO GENERATE STRUCTURE-SPECIFIC PEPTIDES

### Related Applications

5

This application claims the benefit of U.S. Provisional Applications Serial No. 60/211,892, filed June 14, 2000, and Serial No. 60/213,387, filed June 23, 2000. The entire content of each of these provisional applications is hereby incorporated by reference in its entirety.

10

### Field of the Invention

This invention relates to the use of a coiled-coil structural scaffold to generate structure-specific peptides, including synthetic peptides derived from naturally occurring microbial and non-microbial protein antigens. The structure of the synthetic peptides utilizes a scaffold of heptad repeat units into which epitopes derived from coiled-coil regions of native proteins are spliced. The resulting peptide has a more stable coiled-coil structure, hence improving presentation of the epitopes in a helical conformation.

20

### References

U.S. Patent No. 6,075,181.

U.S. Patent No. 6,150,584.

25

WO 96/11944, published April 25, 1996.

WO 01/00010, published January 4, 2001.

30

Alexander J. E., et al. *Infect. Immun.* **62**: 5693-5688 (1994).

-2-

Baquero, F., et al. *J. Antimicrob. Chemother.* **28** (Suppl. C): 31-38 (1991).

Beuvery, E. C., et al. *Infect. Immun.* **37**: 15-22 (1982).

5 Briles D.E. et al. *Infect. Immun.* **57**(5): 1457-1464 (1989).

Briles, D.E. et al. Pneumococcal diversity: considerations for new vaccine strategies with emphasis on pneumococcal surface protein A (PspA). *Clin. Microbiol. Rev.* **11**(4):645-657 (1998).

10

Briles, D. E., et al. In *Effects of microbes on the immune system* (Cunningham, M. W., Flinami, R.S. Eds) Philadelphia: LippinconRaven: pp 263-280 (1999).

Briles, D. E., et al. *Vaccine* **18**: 1707-1711 (2000).

15

Brooks-Walter, A., et al. *Infect. Immun.* **67**: 6533-6542 (1999).

Canvin, J. R., et al. *J. Infect. Dis.* **172**: 119-123 (1995).

20

Chang K, Pastan I. "Molecular cloning and expression of a cDNA encoding a protein detected by the K1 antibody from an ovarian carcinoma (OVCAR-3) cell line", *Int J Cancer.* **57**(1):90-7 (1994).

Cowan, M. J., et al. *Pediatrics* **62**: 721-727 (1978).

25

Crain M. J., et al. *Microb. Pathog.* **21**: 265-275 (1996).

Davis,S., Aldrich,T.H., Jones,P.F., Acheson,A., Compton,D.L., Jain,V., Ryan,T.E., Bruno,J., Radziejewski,C., Maisonpierre,P.C.and Yancopoulos,G.D.  
30 Isolation of angiopoietin-1, a ligand for the TIE2 receptor, by secretion-trap expression cloning. *Cell* **87** (7): 1161-1169 (1996).

De, B. K., et al. *Pathobiology* **67**: 115-122 (1999).

35

Henrichsen, J., *J. Clin. Microbiol.* **33**: 2759-2762 (1995).

Houston, M.E. et al. *Biochemistry* **35**: 10041-10050 (1996).

40

Kalin, M. Pneumococcal serotypes and their clinical relevance. *Thorax* **53**: 159-162 (1998).

Kapanadze B, Kashuba V, Baranova A, Rasool O, van Everdink W, Liu Y, Syomov A, Corcoran M, Poltaraus A, Brodyansky V, Syomova N, Kazakov A,

-3-

- Ibbotson R, van den Berg A, Gizatullin R, Fedorova L, Sulimova G, Zelenin A, Deaven L, Lehrach H, Grander D, Buys C, Oscier D, Zabarovsky ER, Yankovsky N. A cosmid and cDNA fine physical map of a human chromosome 13q14 region frequently lost in B-cell chronic lymphocytic leukemia and  
5 identification of a new putative tumor suppressor gene, Leu5. *FEBS Lett.* **426(2)**:266-70 (1998).
- Lancet, *Lancet* **2**: 699-701 (1985).
- 10 Lawrence, M. C., et al. *Structure* **6**: 1553-1661 (1998).
- Li Y, Kang J, Friedman J, Tarassishin L, Ye J, Kovalenko A, Wallach D, Horwitz MS. Identification of a cell protein (FIP-3) as a modulator of NF-kappaB activity and as a target of an adenovirus inhibitor of tumor necrosis factor  
15 alpha-induced apoptosis. *Proc Natl Acad Sci U S A.* **96(3)**:1042-7 (1999).
- Li,L., Li,X., Francke,U. and Cohen,S.N. The TSG101 tumor susceptibility gene is located in chromosome 11 band p15 and is mutated in human breast cancer. *Cell* **88 (1)**: 143-154 (1997).
- 20 McDaniel. L. S., et al. *Microb. Pathog.* **17**: 323-337 (1994).
- McDaniel. L.S., et al. *Infect. Immun.* **59**: 222-228 (1991).
- 25 Michon F., et al. *Vaccine* **16**: 1732-41 (1998).
- Miranda C, Minoletti F, Greco A, Sozzi G, Pierotti MA. "Refined localization of the human TPR gene to chromosome 1q25 by in situ hybridization", *Genomics* **23(3)**:714-5 (1994).
- 30 Nabors, G. S., et al. *Vaccine* **18**: 1743-1754 (2000).
- Nakashima,M., Sonoda,K. and Watanabe,T. Inhibition of cell growth and induction of apoptotic cell death by the human tumor-associated antigen RCAS1.  
35 *Nat. Med.* **5 (8)**: 938-942 (1999).
- Remington's Pharmaceutical Sciences*, Mack Publishing Company, Philadelphia PA 18<sup>th</sup> ed. (1990).
- 40 Robbins, J. B., et al. *J. Infect. Dis.* **148**: 1136-1159 (1983).
- Shapiro, E. D., et al. *N. Engl. J. Med.* **325**: 1453-1460 (1991).

-4-

Shelly, M. A., et al. *Infect. Immun.* **65**: 242-247 (1997).

Singh et al., "Advances in vaccine adjuvants", *Nat. Biotechnol.* **17(11)**:1075-81 (1999).

5

Spika, J. S., et al. *J. Infect. Dis.* **163**: 1273-1278 (1991).

Takala. A. K., et al. *J. Infect. Dis.* **164**: 982-986 (1991).

10

Talkington, D. F., et al. *Infect. Immun.* **59**: 1285-1289 (1991).

Talkington, D. F., et al. *Microb. Pathog.* **21**: 17-22 (1996).

Tart, R. C., et al. *J. Infect. Immun.* **173**: 380-386 (1996).

15

Tripet, B. et al., *J. Mol. Biol.* **300**: 377-402 (2000).

Wagschal, K. et al., *Protein Science* **8**: 2312-2329 (1999).

20

Winter et al., *Nature* **349**: 293 (1991).

Wolf, E. et al. MultiCoil: A program for predicting two- and three-stranded coiled coils". *Protein Science* **6**: 1179-1189 (1997).

25

World Health Organization. *WHO* 9 (1995).

Yamada et al., "Identification of a novel autoantigen UACA in patients with panuveitis", *Biochem. Biophys. Res. Commun.* **280(4)**:1169-76 (2001).

30

All of the above publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if the disclosure of each individual publication, patent application or patent was specifically and individually indicated to be incorporated by reference in its entirety.

35

#### Background of the Invention

*Streptococcus pneumoniae* is an important pathogen causing life-threatening invasive diseases such as pneumonia, meningitis and bacteraemia. It is estimated

-5-

that more than 1 million cases of pneumococcal pneumonia are diagnosed each year in the United States with 0.7% of these infections being fatal (Lancet, 1985). In developing countries, it is estimated that 4 million deaths are due to pneumonia, with *S. pneumoniae* infections accounting for 70% of the deaths (WHO, 1995). *S. pneumoniae* also causes less serious diseases such as otitis media and sinusitis which, due to their prevalence, are a significant burden on health care systems. The high morbidity and mortality rate associated with pneumococcal infections is exacerbated by the rate at which the organism acquires resistance to multiple antibiotics (Spika et al., 1991; Baquero et al., 1991). Thus, there is an unmet need for effective treatments for pneumococcal infections.

The current vaccine utilized for prevention of pneumococcal infection in humans is based on purified capsular polysaccharides. The design of a capsular polysaccharide vaccine is complicated by the fact that there are 90 different capsular types and the protection elicited by the capsule is type specific (Henrichson, 1995). These problems are somewhat mitigated by the fact that certain capsules are more commonly associated with human disease than others and which ultimately led to their inclusion into the current 23 valent vaccine (Robbins et al., 1983). However, capsular polysaccharides, like most polymeric compounds possessing multiple repeating units, are inefficient in stimulating immune responses and subclass switching to IgG (Beuvery et al., 1982). Not surprisingly, the vaccine is only 60% effective in preventing fatal pneumococcal bacteraemia in the elderly (Shapiro et al., 1991) and is unable to elicit adequate antibody responses in children under the age of 2 (Cowan et al., 1978). The problem of poor immunogenicity in children is being addressed by conjugation of the polysaccharides to protein carriers such as diphtheria and tetanus toxoids (Shelly et al., 1997). In a recent clinical trial, a seven valent conjugate vaccine



-6-

was shown to be immunogenic in children and elicited protection against invasive pneumococcal infection.

5 However, a separate vaccine will be required for adults because clinically important infections are caused by different capsular types than is the case for children. Coverage in the developing world is expected to be as low as 52% owing to geographical variation of pneumococcal strains. In addition, it is anticipated that the conjugate vaccines may be too expensive for the developing world. Thus, there is an urgent need for alternative approaches to the  
10 development of pneumococcal vaccines.

The drawbacks associated with the polysaccharide vaccine have prompted interest in the possibility of developing vaccines based on pneumococcal protein antigens. Pneumococcal proteins under consideration have included the toxins,  
15 pneumolysin, autolysin, and the surface proteins. The surface proteins which have been used as vaccines to date include pneumococcal surface adhesion A (PsaA), pneumococcal surface protein A (PspA) and pneumococcal surface protein C (PspC). The surface proteins are shown in Figure 1.

20 Pneumolysin is a thiol cytolysin that is produced by virtually all clinical isolates of *S. pneumoniae*. When mice were immunized with a genetically engineered toxoid version of pneumolysin (pneumolysoid) absorbed onto alum, the survival rate and the time of death of non-survivors (survival time) was significantly greater than that of controls. However, significant differences in the  
25 degree of protection were noted between different strains (Alexander et al., 1994). Pneumolysoid is currently being investigated as a carrier protein for pneumococcal capsular polysaccharides (Michon et al., 1998). In general, a higher response to

-7-

the capsular polysaccharides was observed when they are formulated as conjugates of pneumolysin as compared to tetanus toxoid conjugates. Immunization with autolysin has also shown to be protective in a sepsis model (Canvin et al., 1995). However, antibodies to autolysin appear to exert their effect in part because they  
5 prevent the autolysin dependent release of pneumolysin.

All *Streptococcus pneumoniae* isolates investigated to date express the lipoprotein pneumococcal surface adhesin A (PsaA). PsaA is 37 kDa in molecular weight and is thought to be genetically conserved. The function of PsaA is that of  
10 a Zn binding membrane transport protein (Lawrence, et al. 1998). Recent studies have shown that immunization of mice with native PsaA elicited anti-PsaA antibodies that protected mice from challenge with virulent strains of *S. pneumoniae* (Talkington et al., 1996; De et al., 1999). In a mouse carriage model, PsaA has been shown to be one of the best pneumococcal surface antigens  
15 at eliciting protection against carriage. However, PsaA is not as effective as PspA in eliciting protection in a murine sepsis model or murine pulmonary infection model (Briles et al., 2000).

Pneumococcal surface protein A has been found on all strains of *S. pneumoniae*  
20 *pneumoniae* studied to date and is required for full virulence (Crain et al., 1990). While PspA has a strain-dependent molecular weight ranging from 67 to 99 kDa, all proteins consist of four distinct domains: an N-terminal highly charged domain, a proline rich domain, a repeat domain comprising 10 highly conserved  
20 amino acid repeats, and a short hydrophobic domain (Figure 2). The repeat  
25 region is responsible for attachment of the protein to the cell surface of *S. pneumoniae* by non-covalent binding to choline residues.



-8-

The amino acid sequence of the Rx1 strain PspA is included herein as SEQ ID NO:1. Analysis of the sequence of the N-terminus of this PspA (residues 1-303) indicated that the sequence adopts a coiled-coil structure. The coiled-coil structure consists of two amphipathic  $\alpha$ -helices wrapped around each other with a left handed supertwist (Figure 3). The coiled-coil is characterized by a heptad repeat (abcdefg), in which the a and d positions are typically occupied by hydrophobic amino acids. These residues are aligned such that they create a hydrophobic face that is responsible for the stability of this structure.

The N-terminal sequence of PspA contains three prolines interspersed throughout the sequence, which are believed to create small breaks in the otherwise continuous coiled-coil. It is proposed from hydrodynamic characterization of the protein and from computer modeling that PspA forms a coiled-coil by folding back onto itself at one of the three proline regions. It is thought that the coiled-coil region extends from the cell wall and possibly protrudes outside the capsule.

The exact function of PspA is not known. It has been shown to retard to clearance of pneumococci from the blood (McDaniel et al., 1987). It seems likely that the major effect of PspA on virulence may be due to interference with complement fixation. In addition, it is thought that the highly positively charged nature of the coiled-coil region may interact with the negatively charged capsule. This interaction may stabilize the capsular structure.

*Streptococcus pneumoniae* has another surface protein called PspC that has a similar domain structure as PspA, but has a larger coiled-coil region and is larger in overall size (Figure 2). The N-terminal 150 amino acids of the helical

-9-

domain are highly variable in size and sequence. Adjacent to this hypervariable region are two repeating sequences that vary in size from 101 to 205 amino acids in length. In the larger PspC proteins, there exists a helical region similar to the C-terminal portion of the coiled-coil region found in PspA. A comparison of PspC and PspA proteins is shown in Figure 2. Evidence now exists that most strains of *S. pneumoniae* produce both PspA and PspC (Brooks-Walter et al., 1999; Crain et al., 1996). The exact function of PspC is not known, but it is believed to function in an analogous fashion as PspA.

#### 10 Summary of the Invention

This invention relates to the unexpected discovery that the characteristic heptad repeat (abcdefg)<sub>n</sub> of coiled-coil proteins, where the a and d positions are typically occupied by hydrophobic amino acids, can be used as a template to lock epitopes from coiled-coil regions or potential coiled-coil regions of native proteins into a stable coiled-coil structure to form novel synthetic peptides based on native proteins. Such peptides may be used for immunization. Amino acid residues from corresponding positions of native coiled-coil proteins are spliced into the solvent exposed b, c, e, f and g positions, while hydrophobic amino acids (in particular, Ile and Leu) occupy the a and d positions. The resulting novel peptides maintain a stable coiled-coil structure in aqueous solution, and present immunogenic epitopes characteristic of the native proteins.

Accordingly, one aspect of the invention provides for synthetic peptides of the following formula (formula I):



-10-

wherein A is Ile, Leu, Val or a derivative thereof, D is Leu, Ile, Val or a derivative thereof, each X is an amino acid residue or derivative thereof which corresponds to an amino acid residue of an epitope of a native coiled-coil protein to form a set of X residues, and n is an integer equal to or greater than 1.

5 Preferably, A is Ile and D is Leu in every set of the (AXXDXXX) repeat. Preferred values for n are 2 to 100. In a preferred embodiment, n is 2 to 10, most preferably about 3 to 6.

10 Preferred sets of X residues are amino acids that are solvent exposed in the coiled-coil region of the native protein. Each set of X amino acid residues may be selected and/or combined independently. In other words, they may come from the same epitope of the same protein, different epitopes of the same protein, epitopes of different proteins, or the like.

15 The peptides may further comprise additional amino acids, such as the amino acid residues CNleG-, at the N-terminus of the peptide. The peptides may also comprise additional amino acids at the C-terminus. The additional amino acids may or may not be arranged in a sequence which has the propensity to form coiled-coils, and the number of additional amino acids at either terminus is preferably less than 100. The number of additional amino acids at either terminus is more preferably less than 50, yet more preferably less than 30, still more preferably less than 20, and most preferably less than 10.

25 This invention may particularly be used to create novel synthetic peptides relating to microbial proteins which exist as coiled-coil structures in the native state. In particular, amino acid residues from epitopes of microbial surface proteins, most particularly *S. pneumoniae* surface proteins A and C, may be used

-11-

in the novel peptides. Many microbial surface proteins have similar architecture, existing as long, free-standing coiled-coils.

The X residues may come from consensus sequences of native coiled-coil proteins, for example PspA and PspC. A preferred consensus sequence of PspA is  
5 EELX<sub>1</sub>X<sub>2</sub>KIDELDX<sub>3</sub>EIAX<sub>4</sub>LEKX<sub>5</sub> (SEQ ID NO:5). Preferably, X<sub>1</sub> is S, Q, N or D; X<sub>2</sub> is D, N or K; X<sub>3</sub> is A or N; X<sub>4</sub> is K, E or D; and X<sub>5</sub> is N, D or E.

To increase the stability of the coiled-coil conformation, the amino acid  
10 residues at the e and g positions, if not part of the epitope of interest, may optionally be replaced with charged amino acids (e.g., Asp, Glu, Lys, Arg and His). The replacement should be performed such that the e position in one peptide strand and the g' position in the "complementary strand" are occupied by amino acids of opposite charges, for example Asp and Lys<sup>1</sup>. Similarly, the residues at  
15 the g position of one strand and the e' position of the complementary strand can be replaced by residues of opposite charges.

The residues may also be replaced by charged amino acids such that a salt bridge can form between two residues which are 3 or 4 residues apart in the same  
20 strand. Alternatively, a charged amino acid may stabilize the coiled-coil structure by interacting with an oppositely-charged residue in a stabilizing strand.

In another aspect, the invention provides methods of making peptides of the formula I comprising the steps of selecting an antigenic epitope derived from a

---

<sup>1</sup>For example, chain 1 and chain 2 in Figure 3 are "complementary strands". As illustrated in Figure 3, the e position in chain 1 and the g' position in chain 2 are aligned. Therefore, if amino acid residues with opposite charge occupy these two positions, they can form a salt bridge between their side chains, which increases the stability of the coiled-coil structure.

-12-

coiled-coil, determining which amino acid residues are solvent exposed, and inserting said solvent exposed amino acid residues into the X positions of formula I. In particular, the selection of epitopes may be performed using a computer algorithm. More than one set of epitopic amino acids derived from the same  
5 protein or different proteins may be used. Furthermore, the sets of epitopic amino acids may be from different strains and/or species of microorganism when epitopes of a microbial coiled-coil protein is used.

In a further aspect, the invention provides compositions useful to stimulate  
10 an immune response in an animal wherein said compositions comprise a peptide of formula I. When more than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the composition is useful to stimulate an immune response to more than one strain and/or species of microorganism. The animal  
15 may be an avian, a mammal (including a human), or any animal capable of an immune response.

Antibodies raised by administration of a peptide of formula I are also provided as an aspect of the invention. Such antibodies may be polyclonal or  
20 monoclonal. Pharmaceutical compositions comprising such antibodies are also provided. When more than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the antibody will bind to more than one strain and/or species of microorganism.

25

In yet a further aspect, the invention provides compositions useful as vaccines wherein said compositions comprise a peptide of formula I. When more

-13-

than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the composition is useful to provide cross protection to more than one strain and/or species of microorganism.

5

In still a further aspect, the invention provides methods of preventing a microbial infection comprising administering to a mammal susceptible to said infection a peptide of formula I. When more than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the composition is useful to prevent infection by several strains and/or species of microorganism.

10

In yet another aspect, the invention provides a method of treating or preventing microbial infection in an animal susceptible to or suffering from such infection, comprising administering to said animal an effective amount of an antibody to a microbial protein, wherein said antibody is produced by administering a peptide of formula I to an animal. In particular, such administration may be used to provide passive immunization and/or prevent or alleviate symptoms of infection in said animal. When more than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the method may treat or prevent infection by several strains and/or species of microorganism.

15

20

In still another aspect, the invention provides a method of determining the presence of a particular microorganism in a sample comprising contacting the sample with an antibody to a peptide of formula I which peptide comprises

25



-14-

epitopes from the particular microorganism, and determining whether said antibody binds to a component of said sample. The sample may be a biological sample. The method may be used to determine the causative agent of a microbial infection. When more than one set of epitopic amino acids is used in the peptide  
5 of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism, the method may be used to simultaneously detect the presence of several strains and/or species of microorganism in the sample.

In a further still aspect, the invention provides a method for providing  
10 multiple epitopes for exposure to the immune system of an animal comprising administering to the animal a peptide of formula I wherein said peptide epitopes are from different proteins. In a preferred embodiment, the peptide epitopes are from different proteins of a single microorganism or from different strains and/or species of microorganisms.

15 In still a further aspect, the invention provides a method for determining the presence of antibodies to a microbial protein in a biological sample comprising contacting the sample with a peptide of formula I which peptide comprises at least one epitope from the microbial protein and determining binding of antibodies in  
20 the sample to the peptide. The method may be used to determine whether an animal has been exposed to a microorganism.

#### Brief Description of the Drawings

25 Figure 1 is a hypothetical representation of the pneumococcal surface depicting several antigens present on the surface. The coiled-coil regions from PspA and PspC are believed to project away from cell wall.

-15-

Figure 2 is a diagrammatic representation of the PspA and PspC molecules highlighting their regions of homology. Each protein consists of 4 domains, a coiled-coil domain, a proline rich region, a choline binding region and a hydrophobic tail. High sequence homology exists in the region 2 sections of the coiled-coil domains between PspA and PspC.

Figure 3 depicts the cross-sectional view of a coiled-coil structure consisting of two peptide chains. The 7 positions of the heptad repeat in chain 1 are designated a, b, c, d, e, f and g, while the 7 positions of the heptad repeat in chain 2 are designated a', b', c', d', e', f' and g'. The open arrows indicate hydrophobic interactions between a and a', and d and d', respectively.

Figure 4 shows the steps of combining the epitopes of several clinically relevant strains of *S. pneumoniae* into a coiled-coil template to make an immunogenic peptide.

Figure 5 shows the CD spectrum of the peptide of Figure 4 in the presence and absence of TFE.

Figure 6 shows a region of PspA which is highly conserved. A consensus sequence which is selected such that it conforms with formula I is also shown.

Figure 7 shows the CD spectrum of peptide CVX0270, which contains a portion of the consensus sequence shown in Figure 6.

Figure 8 shows the thermal denaturation profile of peptide CVX0270.

-16-

Detailed Description of the Invention

This invention relates to the use of a coiled-coil structural scaffold to generate structure-specific peptides, including synthetic peptides derived from naturally occurring protein antigens. The structure of the synthetic peptides utilizes a scaffold of heptad repeat units into which epitopes of native coiled-coil proteins are spliced.

In particular, the synthetic peptides are based on microbial proteins, especially surface proteins, which occur naturally in the coiled-coil form, such as pneumococcal surface proteins A and C. The synthetic peptides are immunogenic and can be used to elicit an immune response in an animal. Accordingly, they are useful as vaccines or to stimulate antibody production or cell-mediated immunity which recognizes the naturally occurring protein.

Prior to describing the invention in further detail, the terms used in this application are defined as follows unless otherwise indicated.

Definitions

A “native protein” is a protein which exists in nature.

A “coiled-coil protein” is a protein which, at least in part, forms a coiled-coil. A coiled-coil protein may have other conformational structures in addition to the coiled-coil. Encompassed within the term coiled-coil proteins are the proteins which have been shown to assume a coiled-coil structure and the proteins which are predicted to form coiled-coils by using a computer algorithm.

-17-

A “microbial protein” is a protein derived from a microorganism such as a bacterium, archaeobacterium, fungus, virus, protozoan, parasite, alga, slime mold, or prion.

5           A “peptide” as used herein is a peptide or a protein. In other words, a peptide referred to in this application may contain any number of amino acids.

A “peptide of formula I” is a peptide which comprises a synthetic fragment, which fragment consists of formula I.

10

An “epitope” is a part of a protein or peptide which is an antigenic determinant.

15           A “derivative” of an amino acid is a non-naturally occurring amino acid residue or a chemically modified amino acid. Amino acid derivatives may be used to increase the half life of the peptide in serum or tissue, or to increase antigenicity of the peptide.

20           Non-naturally occurring amino acids may be but not limited to D-isomers, norleucine, 4-amino butyric acid, aminoisobutyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid and 2-thienyl alanine.

25           A chemically modified amino acid is an amino acid with a side chain modification. For example, the amino group of lysine may be modified by alkylation with an aldehyde followed by reduction with  $\text{NaBH}_4$ ; amidation with

-18-

methylacetimidate; acylation with acetic anhydride; carbamoylation with cyanate; trinitrobenzylation with 2,4,6,-trinitrobenzene sulphonic acid (TNBS); acylation with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation with pridoxal-5'-phosphate followed by reduction with  $\text{NaBH}_4$ .

5

The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

10

The carboxyl group may be modified by carbodiimide activation via O-acylisourea formation followed by subsequent derivatization, for example, to a corresponding amide.

15

The sulfhydryl group may be modified by carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulfide with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted maleimide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; or carbamoylation with cyanate at an alkaline pH.

20

Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

25

-19-

Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or diethylpyrocarbonate.

5 A "peptide" may be a peptide containing naturally existing amino acids or amino acid derivatives. A peptide may also contain cross links between different portions of the peptide. For example, coiled-coil structures are often composed of two strands or three strands. It is preferable that the different strands of the coiled-coil are linked by disulfide bonds, which stabilize the coiled-coil structure.  
10 More preferably, the peptide contains a lactam bridge between the sidechains of lysine and glutamic acid which are spaced 3 or 4 residues apart (Houston et al., 1996). These lactam bridges are incorporated preferably at the N- and C-termini of the peptide sequence and not in regions of the sequence where the epitope is being displayed.

15 A "solvent exposed" or "solvent accessible" amino acid residue is an amino acid residue of a protein or peptide which is exposed to the solvent when the protein or peptide exists in a solution. The solution is preferably an aqueous solution, more preferably a physiologically compatible solution, such as blood,  
20 lymphatic fluid or the benign buffer. In particular, "solvent exposed" amino acid residues refer to the residues at the b, c, e, f or g positions of a native epitope which forms or is predicted to form a coiled-coil.

A benign buffer is a phosphate buffered solution. The "benign buffer" as  
25 used in this application is 50 mM  $\text{KH}_2\text{PO}_4$ , 50 mM KCl, pH 7.0.



-20-

An "antibody" is a protein molecule that reacts with a specific antigen and belongs to one of five distinct classes based on structural properties: IgA, IgD, IgE, IgG and IgM.

5           An "immune response" is the development in the host of a cellular and/or antibody-mediated immune response to a composition or vaccine of interest. Such a response may consist of the production of one or more of the following: antibodies, B cells, helper T cells, suppressor T cells, and/or cytotoxic T cells directed specifically to an antigen or antigens included in the composition or  
10       vaccine of interest.

A "vaccine" is a molecule which is capable of eliciting in an animal an immune response which prevents, partially or completely, infection by a pathogen.

15           "Elicit" or "stimulate" an immune response is to cause an immune response by exposing an immune system to an immunogen.

An "immunogen" is a molecule which is capable of eliciting an immune response in an animal.  
20

"Prevent a microbial infection" is to prevent, completely or partially, the development of a microbial infection.

"Treat a microbial infection" is to reduce, completely or partially, the  
25       symptoms of a microbial infection after the onset of the microbial infection.

-21-

A “sample” is an aliquot or a representative portion of a substance, material, or population. For example, a sample may be a sample of water, sewage, oil, sand, blood, biological tissue, urine or feces.

5           A “biological sample” is a sample collected from a biological subject, such as an animal, plant or microorganism.

10           An “effective amount” is an amount which is sufficient to achieve the intended purposes. For example, an effective amount of a vaccine is an amount of the vaccine sufficient to elicit an immune response in the recipient of the vaccine to protect the recipient from contracting the target disease or to prevent or alleviate medical conditions associated with the disease.

The synthetic coiled-coil peptide

15

It has been shown previously that PspA is highly variable among different strains of *S. pneumoniae*. Mapping studies using monoclonal antibodies raised against PspA indicate that the major cross-reactive epitopes are found in the last 100 amino acids of the coiled-coil domain (residues 192-270 of SEQ ID NO:1) (McDaniel et al., 1994). Based on the sequence of this region, it has been possible to group PspA proteins into 6 clades, which comprise three families. Animal studies have shown that immunization with PspA elicits an immune response which is not only cross-reactive to other strains in the same PspA family, but that this cross-reactivity extends between families (Briles, et al. 1999; McDaniel et al., 1991; Tart et al., 1996). In addition, immunization with purified PspC fragments generates antibodies that are protective against sepsis and cross react with PspA (Brooks-Walter et al., 1999). The cross-protection results suggest that despite the

20

25

-22-

variation among the coiled-coil domain for PspA and PspC, there must be conserved epitopes.

The first evidence that immunity to PspA might be protective was the finding that PspA specific monoclonal antibodies (MAbs) protected mice from fatal sepsis. Subsequently, it was shown that passive protection with IgG or IgM monoclonal antibodies to PspA from strain Rx1 protected mice from death following ip or iv challenge with  $10^5$  times the  $LD_{50}$  of *S. pneumoniae* strain WU2 (Briles, et al. 1989). The MAbs were shown to clear the pneumococci from the blood of the challenged mice by a complement-mediated mechanism. Active immunization studies utilized a 27 kDa fragment of PspA from *S. pneumoniae* strain Rx1 (residues 1-245 of SEQ ID NO:1) which contain most of the helical domain of the protein. Immunization of CBA/N mice with 5  $\mu$ g of this fragment in combination with the adjuvant FCA followed by a booster dose in PBS were protected from an iv challenge of 300 cfu (30 times  $LD_{50}$ ) of strain WU2 (Talkington et al., 1991). In more recent studies when mice received 1 or 5  $\mu$ g of a fragment comprising residues 1-303 with no adjuvant, it elicited protection against fatal infection following iv challenge with 480 cfu of the more virulent strain A66.1 (Briles et al. 1998). Passive protection studies in mice with immune sera from immunized mice or rabbits can protect against fatal sepsis from challenge doses of 10 to 100 times  $LD_{50}$  of the challenge strain.

While PspA can be divided into numerous serological types, it has been shown to be a highly cross-reactive protein. Sera from a single rabbit immunized with the helical portion of PspA from strain Rx1 can recognize PspA from all pneumococci in Western blots. When CBA/N mice were challenged with 14 different *S. pneumoniae* strains ( $\sim 200 \times LD_{50}$  dose) following immunization with

-23-

PspA from strains D30, WU2 and BG9739, each PspA exhibited cross-protection to most challenge strains. The helical domain of PspA from the strain Rx1 (residues 1-314 of SEQ ID NO:1) was recently evaluated for safety and immunogenicity in human volunteers (Nabors et al., 2000). When patients were immunized with 5 to 125 mg of this fragment absorbed onto aluminum hydroxide, high levels of circulating antibodies to the Rx1 fragment were elicited as well as antibodies reactive to heterologous PspA molecules.

The PspA sequence from *S. pneumoniae* strain Rx1 was analyzed using the Peptools program from Biotoools Inc. The Peptools structure prediction algorithm is based on identifying known protein-folding motifs combined with consensus prediction based on the results from four prediction algorithms. The Peptools program also predicts a highly helical region that spans over 300 amino acids for the N-terminal portion of PspA. Peptools program is also capable of predicting which sites of proteins are antigenic by measuring hydrophilicity, solvent accessibility, flexibility, and turn propensity of various sequences, and in addition, matches sequences against a database of known B-cell epitopes. Peptools identified a number of potential B-cell epitopes in the coiled-coil domain which are listed in Appendix 1.

The folding of PspA and PspC (or other proteins which exist as coiled-coils) into a coiled-coil results in certain residues being solvent accessible and certain residues which form the hydrophobic core being solvent inaccessible. The solvent accessible amino acids are located at positions *b*, *c*, *e*, *f* and *g* of the heptad repeat while amino acids *a* and *d* are buried. Thus, residues at solvent accessible positions must be responsible for mediating an immune response, e.g., antibody production. The epitopes formed by the coiled-coil region must be discontinuous

-24-

since the *a* and *d* residues are not exposed. Thus, the epitopes from these proteins must be derived from the solvent accessible positions *b*, *c*, *e*, *f* and *g*.

We discovered that the stability of the hydrophobic core of a coiled-coil protein may be maximized by incorporation of Ile residues at *a* positions and Leu residues at *d* positions of the heptad repeat. Accordingly, the present invention provides a synthetic peptide comprising formula I:



wherein

A is Ile or a derivative thereof;

D is Leu or a derivative thereof;

each X is an amino acid residue or derivative thereof which corresponds to an amino acid residue of an epitope of a native coiled-coil protein;

the X residues in each (AXXDXXX) repeat form a set of X residues; and

*n* is equal to or greater than 1.

While the preferred embodiment has Ile at the *a* position and Leu at the *d* position, also contemplated in the present invention are peptides containing Val, Ile or Leu at the *a* or *d* positions. Thus, the amino acid at the *a* position may be Val, Ile or Leu, and the amino acid at the *d* position may, independently, also be Val, Ile or Leu.

Although the coiled regions of PspA and PspC are quite long, synthetic coiled-coil forming sequences may be shorter. It has been shown that if lactam bridges are included, a synthetic coiled-coil may be as short as 14 residues

-25-

(Houston et al., 1996). In the absence of lactam bridges, we have found that helical peptides containing 21 amino acids were stable. In addition, linking of the individual helices together by an intermolecular disulfide bond further stabilizes these structures. The small coiled-coils can serve as scaffolds into which epitopes  
5 from native coiled-coil proteins, such as PspA and PspC, are spliced. A potential scaffold has the following sequence:



wherein Nle is norleucine, an isomer of leucine containing an unbranched side chain. Residues that make up epitopes from native coiled-coil proteins such as  
10 PspA and PspC (residues *b*, *c*, *e*, *f* and *g*) would be inserted into the X positions. The epitopes could potentially be determined by using algorithms and bioinformatics programs such as Peptools. Epitopes could also be mapped by monoclonal antibodies or by analysis of polyclonal antisera raised against native proteins on a Biacore instrument. Epitopes from different native proteins such as  
15 from different strains of *S. pneumoniae* could be spliced together to yield novel sequences. The peptide may be synthesized by peptide synthesis methods established in the art, or an expression vector encoding the peptide sequence of interest can be constructed and used to express the peptide. The I and L residues in this scaffold may, independently, be replaced by any amino acid selected from  
20 I, L or V.

The amino acids at the *b*, *c*, *e*, *f*, and *g* positions in the present invention are the solvent exposed amino acids from natural coiled-coils of the particular protein of interest. Therefore, the present invention is distinct from WO  
25 01/00010, which is directed to displaying solvent exposed amino acids from a non-coiled-coil protein in a coiled-coil template. As a result, a non-coiled-coil epitope is converted to a coiled-coil conformation, and it is possible that antibodies raised



-26-

against the coiled-coil conformation will not recognize the original, non-coiled-coil epitope. In contrast, a peptide made according to formula I of the present invention contains at the b, c, e, f, and g positions the amino acids from the b, c, e, f, and g positions of a coiled-coil protein. The resulting peptide, while being  
5 more immunogenic, will elicit the same immune response as the original epitope with respect to specificity.

The present invention is also distinct from WO 96/11944, which is directed to embedding one peptide in another. Briefly, WO 96/11944 discloses that one  
10 can enhance the immunogenicity of a peptide by flanking the peptide with amino acids which have a strong propensity to form the same conformation as the peptide of interest, thereby stabilizing the conformation of the peptide. For example, a peptide which may form a coiled-coil can be embedded in the amino acid sequences of GCN4, which forms relatively stable coiled-coils. The resulting  
15 chimeric peptide had a high helical content when dissolved in the  $\alpha$ -helix inducing solvent trifluoroethanol, but its helical content under aqueous conditions was low. The present invention, on the other hand, does not embed a peptide fragment in another amino acid sequence. Instead, the amino acids at the a and d positions of the peptide are specifically chosen and replaced where appropriate in order to  
20 increase the stability of the coiled-coil conformation, and the resulting peptide is typically helical even in aqueous solutions.

The following example demonstrates the basic features of the present invention. As mentioned previously, the C-terminal portion of the coiled-coil  
25 domain from the Rx1 PspA protein has been shown to contain a number of cross-protective epitopes. The Biotools software identified a number of such epitopes including residues 153-170 and residues 181-198 that will be used to illustrate the

-27-

construction of a coiled-coil immunogen (Example 1). The *a* and *d* position of these epitopes are highlighted in bold text while the solvent accessible amino acids are in italics. These positions are replaced by Ile at *a* positions and Leu at *d* positions of the template. By simply splicing out the solvent exposed residues (italicized) from the potential epitopes and incorporating them into the scaffold,  
5 one is able to create unique polypeptide sequences which are immunogenic. The coiled-coils described in Example 1 will have approximate molecular weights of 7,000 Da and may be large enough to elicit an immune response on their own. Alternatively, these molecules could be coupled to carrier proteins such as tetanus  
10 toxoid or KLH, or attached to liposomes using techniques known to those of skill in the art.

It must be stressed that the residues must be kept in phase such that, for example, a *c* residue from the protein must correspond to an *c* residue in formula  
15 I. Residues from either of the two adjoining sequences can be used to fill any gaps that might arise during the splicing. Since the coiled-coil is composed of two  $\alpha$ -helices, the two helical strands can bear different sequences, thus increasing the number of epitopes in the entire molecule. Through incorporation of a number of epitopes from different clades, the coiled-coil peptides will cross react among  
20 different clades.

By maintaining a constant hydrophobic core, splicing in only solvent exposed residues and mixing epitopes from different proteins, unique sequences are produced. Thus, a formula I peptide may have the amino acids from the *b*, *c*,  
25 *e*, *f* and *g* positions of PspA in one heptad repeat and those from PspC in another repeat. When this peptide is used as a vaccine, immunity against both PspA and PspC is induced, thereby providing double protection against *S. pneumoniae*.

-28-

A particularly appealing approach is to include amino acids from both the surface proteins (such as PspA and PspC) and the surface adhesin protein (PsaA). The adhesin protein is responsible for pathogenesis of *S. pneumoniae* in the nasopharyngeal area while the surface proteins are involved in systemic symptoms. It has been shown that PsaA may be an effective vaccine to prevent nasopharyngeal carriage while PspA is a better vaccine candidate for systemic infections (Briles et al., 2000). Therefore, a peptide comprising epitopes from both PsaA and PspA will provide a defense against both the nasopharyngeal and systemic infections by the bacteria. This can be achieved by splicing the solvent exposed amino acids from the  $\alpha$ -helical region of PsaA into one heptad repeat of a Formula I peptide, and the solvent exposed amino acids from the  $\alpha$ -helical region of PspA into another heptad repeat of the same peptide. Similarly, a hybrid peptide comprising both PspC epitopes and PsaA epitopes can be prepared.

Alternatively, the formula I peptide may contain the solvent exposed amino acids of proteins from different strains of a microorganism or from different microorganisms, and the resulting vaccine can be effective against all the strains or microorganisms used. Similarly, a consensus sequence may be deduced for a certain protein from different strains and used as a vaccine against all these strains (see Example 3).

In order to improve the helical stability of a peptide, the e and g positions of the heptad repeat of formula I may contain glutamic acid and lysine. These residues form salt bridges between e of one strand and g' of the other strand, as well as g of one strand and e' of the other strand. These salt bridges further help to stabilize the coiled-coil conformation. To stabilize individual  $\alpha$ -helices, lysine residues and glutamic acid residues can be introduced at positions 3-4 residues

-29-

apart to facilitate the formation of an intrastrand salt bridge between their sidechains. One can also increase the stability of the coiled-coil by the inclusion of amino acids with high helical propensity such as alanine or aminoisobutyric acid.

5           Alternatively, a stabilizing strand may be used which does not contain epitopes. The sole purpose of the stabilizing strand is to increase the helical content of the strand which contains the epitopes. A stabilizing strand is capable of forming a very stable helix which supports and stabilizes the helical structure formation of the epitope-containing strand. Preferably, the epitope has lysine  
10           and/or glutamic acid at the e or g positions, and the stabilizing strand can provide an amino acid with the opposite charge at the e' or g' positions to form an interstrand salt bridge. An example of a stabilizing strand is:

CNleGGG(EIEALKK)<sub>5</sub>           (SEQ ID NO:2)

15           The structural and conformational integrity of these peptides can be readily determined by circular dichroism (CD) spectroscopy. The peptides useful in the present invention have helical characteristics in aqueous solutions (see Examples 2 and 3). Therefore, the native epitopes can be correctly presented in a physiological environment. In contrast, previously reported synthetic coiled-coil  
20           proteins do not form helices in aqueous solutions (WO 96/11944). Peptides useful in the present invention have a helical content of at least 20% in benign buffer as determined by CD spectroscopy. The helical content is preferably at least 40%, more preferably at least 60% and most preferably at least 80% in benign buffer.

25           The stability of the peptides may be determined by thermal denaturation assays or chemical denaturation assays such as the guanidine chloride denaturation assay. Such assays are known to those skilled in the art. The stability of the

-30-

peptides may also be indirectly assessed by inhibition ELISA, wherein the ability of a peptide to inhibit the interaction between an antigen and an antibody serves as an indicator of how stable the peptide is in assuming the necessary conformation to compete with the antigen for the antibody.

5

Epitopes from other microbial coiled-coil proteins may also be used in the claimed invention. In particular, epitopes from microbial surface proteins which contain coiled-coil structures may be used. Examples of microorganisms whose proteins are useful in the claimed invention include, but are not limited to,

10 *Haemophilus influenza*, and particularly its hypothetical protein HI0500; the Lyme disease spirochete, i.e., *Borrelia burgdorferi*, in particular its surface lipoprotein P27; *Legionella*, especially its hypothetical protein ppIB; *E. coli*, particularly its conserved YigN protein; *Neisseria* spp, and in particular *Neisseria meningitidis* and its putative periplasmic protein; *Moraxella catarrhalis*, especially its proteins

15 UspA1 and UspA2; *Mycoplasma* spp, and in particular, *Mycoplasma pneumoniae* and its hypothetical protein EO7; *Chlamydia* spp, particularly hypothetical protein CT825; *Plasmodium falciparum*, and particularly its hypothetical protein PFB01450 and mature-parasite-infected erythrocyte surface antigen; *Trypanosoma* spp; *Staphylococcus aureus*; *Streptococcus*, particularly Group A and Group B;

20 *Bordetella pertussis*; *Salmonella*; *Streptococcus mutans*; *Cryptococcus neoforms*; *Klebsiella pneumonia*; *Pseudomonas aeruginosa*; viruses; fungi; and the like.

Moreover, other coiled-coil proteins which do not come from a microbial origin may be used in the same fashion. For example, it is common to raise

25 antibodies to a protein for therapeutic purposes or to study the functions of this protein. However, not every protein of interest is sufficiently immunogenic, and it may be difficult to raise antibodies specifically directed to particular coiled-coil

-31-

regions of the native protein. The present invention may be used to optimize antibody preparation for any coiled-coil protein no matter what origin, function or subcellular localization this protein may have. Moreover, the present invention also enables production of antibodies directed to particular coiled-coil regions, by including only the region of interest in the peptide used to raise antibodies.

Examples of non-microbial coiled-coil proteins include, without being limited to, estrogen receptor binding fragment associated gene 9 (Nakashima et al., 1999), leukemia associated protein 5 (Kapanadze et al., 1998), uveal autoantigen (Yamada et al., 2001), angiopoietin precursor (Davis et al., 1996), NF-kappa B essential modulator (Li et al., 1999), Tumor susceptibility gene 101 protein (Li et al., 1997), cytosolic ovarian carcinoma antigen 1 (Chang et al., 1994), TACC 1 (Still, 1999), and TPR (Miranda et al., 1994).

In addition to the microbial and non-microbial proteins listed herein, a skilled artisan can identify other coiled-coil proteins according to established methods in the art. For example, if the amino acid sequence of a protein is available, various computer algorithms such as Multicoil or Peptool can be used to predict if the protein forms a coiled-coil. For a protein which is substantially pure, CD spectroscopy can be used to determine its actual structure as described herein.



-32-

The compositions

A composition comprising the coiled-coil peptides of formula I can be used to elicit an immune response in an animal for at least two purposes. Where the composition acts as a vaccine by eliciting an immune response in the animal, the resulting antibodies or T-cell mediated immunity can protect the animal from a subsequent attack involving the same epitopes (active immunity). Alternatively, the composition can be used to produce antibodies which can be used as a research tool, or administered to a second animal to protect the second animal from a subsequent attack involving the same epitopes (passive immunity).

To augment the immune response elicited, it may be preferable to couple the peptides of formula I, especially the smaller peptides (e.g., those containing one to four heptad repeats), to a carrier protein.

In addition, the coiled-coil peptides of formula I or their conjugates with carrier proteins may be further mixed with adjuvants to elicit an immune response, as adjuvants may increase immunoprotective antibody titers or cell mediated immunity response. Such adjuvants may include, but are not limited to, Freund's complete adjuvant, Freund's incomplete adjuvant, aluminum hydroxide, dimethyldioctadecyl-ammonium bromide, Adjuvax (Alpha-Beta Technology), Inject Alum (Pierce), Monophosphoryl Lipid A (Ribi Immunochem Research), MPL+TDM (Ribi Immunochem Research), Titermax (CytRx), QS21, the CpG sequences (Singh et al., 1999), toxins, toxoids, glycoproteins, lipids, glycolipids, bacterial cell walls, subunits (bacterial or viral), carbohydrate moieties (mono-, di-, tri-, tetra-, oligo- and polysaccharide), various liposome formulations or

-33-

saponins. Combinations of various adjuvants may be used with the antigen to prepare the immunogen formulation.

5 The composition may be administered by various delivery methods including intravascularly, intraperitoneally, intramuscularly, intradermally, subcutaneously, orally, nasally or by inhalation. The composition may further comprise a pharmaceutically acceptable excipient and/or carrier. Such compositions are useful for immunizing any animal which is capable of initiating an immune response, such as primate, rodent, bovine, ovine, caprine, equine, 10 leporine, porcine, canine and avian species. Both domestic and wild animals may be immunized. The exact formulation of the compositions will depend on the particular peptide or peptide-carrier conjugate, the species to be immunized, and the route of administration.

15 The antibodies produced against a coiled-coil protein can be included in a pharmaceutical composition and administered to an animal. The pharmaceutical composition typically comprises a pharmaceutically acceptable carrier, and may include pharmaceutically acceptable excipients. The pharmaceutical composition can be administered intravascularly, intraperitoneally, intramuscularly, 20 intradermally, subcutaneously, orally, nasally or by aerosol inhalation. Preferably the pharmaceutical composition is administered intravascularly, intramuscularly, nasally or by aerosol inhalation.

Also encompassed by the present invention are antibodies, particularly 25 monoclonal antibodies, which are derived from the antibodies produced against a peptide of formula I. In particular, hybridomas can be generated using a peptide of formula I, and recombinant derivative antibodies can be made using these

-34-

hybridomas according to well-known genetic engineering methods (for a review, see Winter et al., 1991). For example, the DNA fragment coding for the variable regions of the monoclonal antibodies can be obtained by polymerase chain reactions (PCR). The PCR primers can be oligonucleotides which are  
5 complementary to the constant regions of the heavy chain or light chain, and the PCR template can be the total cDNA or genomic DNA prepared from the hybridomas. Alternatively, a cDNA library can be prepared from the hybridomas and screened with probes which correspond to the constant regions of immunoglobulin heavy chain or light chain to obtain clones of the heavy chain or  
10 light chain produced by the particular hybridoma.

Subsequently, the DNA fragment for the variable regions can be inserted into an expression vector and joined in frame with the cDNA sequences of a selected constant region. The constant region can be the human constant  
15 sequences to make humanized antibodies, the goat constant sequences to make goat antibodies, the IgE constant sequences to make IgE which recognizes the peptide of formula I, and the like. Thus, antibodies with the same antigen recognition ability but different constant regions can be produced. Of particular interest are humanized antibodies, which can be used as therapeutic agents against a disease  
20 associated with the cognate antigen in humans without eliciting an undesired immune response against the humanized constant region.

Other methods known in the art to humanize antibodies or produce human antibodies can be utilized as well, including but not limited to the xenomouse  
25 technology developed by Abgenix Inc. (U.S. Patent Nos. 6,075,181; 6,150,584) and the methods developed by Biovation, Bioinvent International AB, Protein Design Labs., Applied Molecular Evolution, Inc., ImmGenics Pharmaceuticals

-35-

Inc., Medarex, Inc., Cambridge Antibody Technology, Elan, Eos Biotechnology, MedImmune, MorphoSys or UroGensys Inc. Likewise, other methods known in the art to screen human antibody secreting cells to coiled-coil peptide antigens can also be utilized.

5

The formulation for the composition, comprising either a coiled-coil peptide or an antibody against a coiled-coil peptide, will vary depending on factors such as the administration route, the size and species of the animal to be administered, and the purpose of the administration. Suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*.

10

The following examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of the present invention.

15

-36-

**EXAMPLES**

In the examples below, the following abbreviations have the following meanings. Abbreviations not defined have their generally accepted meanings.

	°C	=	degree Celsius
5	hr	=	hour
	min	=	minute
	μM	=	micromolar
	mM	=	millimolar
	M	=	molar
10	ml	=	milliliter
	μl	=	microliter
	mg	=	milligram
	μg	=	microgram
	rpm	=	revolutions per minute
15	ID	=	inner diameter
	TFE	=	trifluoroethanol
	EDT	=	ethane dithiol
	TFA	=	trifluoroacetic acid
	PBS	=	phosphate buffered saline
20	β-ME	=	β-mercaptoethanol
	DMSO	=	dimethylsulfoxide
	Nle	=	norleucine
	ELISA	=	enzyme linked immunosorbent assay
	HRP	=	horse radish peroxidase
25	TD	=	thymus dependent
	TT	=	tetanus toxoid
	KLH	=	keyhole limpet hemocyanin

-37-

BSA            =        bovine serum albumin

The one letter code and the three letter code for amino acids used throughout this application are listed below:

5

---

A = Ala = Alanine

C = Cys = Cysteine

D = Asp = Aspartic Acid

E = Glu = Glutamic Acid

10

F = Phe = Phenylalanine

G = Gly = Glycine

H = His = Histidine

I = Ile = Isoleucine

K = Lys = Lysine

15

L = Leu = Leucine

M = Met = Methionine

N = Asn = Asparagine

P = Pro = Proline

Q = Gln = Glutamine

20

R = Arg = Arginine

S = Ser = Serine

T = Thr = Threonine

V = Val = Valine

W = Try = Tryptophan

25

Y = Tyr = Tyrosine

---



-38-

## Materials and Methods

### *Peptide synthesis and purification*

5           The peptide analogs were prepared by t-Boc chemistry on an Applied Biosystem 431 A peptide synthesizer. The peptide resin (700 mg) was cleaved with 10 mL of HF containing 10% anisole and 2.5% EDT for 1 h at 0-4°C. After removal of the HF the peptide/resin was transferred to a sintered glass funnel and washed with diethyl ether (2×50 mL) followed by glacial acetic acid (2×50 mL).  
10       The acetic acid solution was then lyophilized. The crude peptide was taken up in 15 mL of water containing 0.05% TFA and 3mL acetic acid. After stirring and sonication, the mixture was transferred to 1.5 mL Eppendorf tubes and centrifuged at 13000 rpm. The supernatant was collected and filtered through a Millex GV 0.22 μm syringe filter. This solution was loaded onto a Zorbax RX-C8 (22.1 mm  
15       ID×250 mm, 5 μm particle size) through a 5 mL injection loop at a flow rate of 3 mL/min. The purification was accomplished by running a linear AB gradient of 0.1% B/min where solvent A is 0.05% TFA in water and solvent B is 0.05% TFA in acetonitrile.

20           A linker and cysteine residue was incorporated to enable the strands of the coiled-coil to be linked together by a disulfide bond. The disulfide-bridged coiled-coils were formed by overnight air oxidation at room temperature of ~ 10 mg/mL peptide in 100 mM NH<sub>4</sub>HCO<sub>3</sub>, pH 8.5. All peptides were characterized by analytical HPLC and electrospray mass spectrometry, and protein concentrations  
25       as well as amino acid composition were determined by amino acid analysis. Peptides were conjugated to the carrier protein tetanus toxoid rising the photoreactive reagent benzoylbenzoic acid which was coupled to the N-terminus of

-39-

the peptide by conventional solid-phase chemistry described above. This photoaffinity probe is stable under the conditions of t-Boc chemistry. A norleucine residue is incorporated into the linker region of the peptide to enable the quantitation of the peptide attached to the carrier protein by means of amino acid analysis. To 10 mg of tetanus toxoid dissolved in 100 mm of  $\text{NH}_4\text{HCO}_3$  pH 8.3 buffer was added 6 mg of peptide and the solution was irradiated for 1 h at a wavelength of 350 nm. To remove any unreacted peptide, the solution was transferred to a 15 mL dialysis cassette and dialyzed against 4 L of 20 mM  $\text{NaH}_2\text{PO}_4$  pH 7.0 buffer. HPLC analysis of the dialysate was used to determine the amount of unreacted peptide present in the irradiated solution. The dialysis was deemed to be complete when residual peptide was less than 1% of the total area under the curve in the HPLC chromatogram.

#### *CD spectroscopy*

15

Circular dichroism (CD) spectroscopy was performed using a Jasco J-500C spectropolarimeter (Jasco, Easton, Maryland) equipped with a Jasco DP-500N data processor. A 10-fold dilution of an  $\sim 500\text{mM}$  stock solution of the disulfide-linked peptide was loaded into a 0.02 cm fused silica cell and ellipticity scanned from 190 to 250 nm. Each disulfide-bridged analog was analyzed by CD spectroscopy under benign conditions (50 mM phosphate, 100 mM KCl, pH 7.0) and also in the presence of 50% TFE in the same buffer. A Lauda water bath (model RMS, Brinkmann Instruments, Rexdale, Ont.) was used to control the temperature of the cell. CD spectra were the average of four scans obtained by collecting data at 0.1 nm intervals from 250 to 190 nm.

25

-40-

*HPLC*

Analytical HPLC was performed on a Beckman System Gold HPLC equipped with a diode array detector. Analyses were performed on a Zorbax SB-C8 column (4.5 mm ID x 150, 3.5  $\mu$ m particle size). Mobile phases were A: deionized purified  
5 water (18 M $\Omega$ /cm) containing 0.05% trifluoroacetic acid and B: acetonitrile (HPLC grade, 99.99%) containing 0.05% TFA. HPLC operating conditions are listed below:

Flow rate	1 mL/min
Injection Volume	20 mL
Detector Wavelength	215 nm
10 Gradient	2.5% B/min
Total run time	20 min
Integration Start Time	5 min
Minimum Area (integration)	25000

15

**Example 1 Construction of a synthetic pneumococcal immunogen**

This Example illustrates the method of splicing the amino acid residues at the b, c, e, f or g positions of the coiled-coil structures of PspA into a synthetic  
20 immunogen. The Biotoools software identified a number of coiled-coil motifs in PspA, including residues 153-170 and 181-198. As shown below, the a and d positions of these motifs are indicated in bold text while the solvent accessible amino acids are italicized. A space is inserted between the heptad repeats for clarity.

25

<b>LEEAEKK</b> <i>ATEAKQK</i> VDA	residues 153-168 of SEQ ID NO:1
<i>AEL</i> <b>ENQ</b> <i>VHRLEQE</i> <b>LKEIDES</b>	residues 181-198 of SEQ ID NO:1

-41-

The solvent exposed residues are incorporated into the coiled-coil template with isoleucine and leucine at the a and d positions, respectively:

5                    **LEEAEKK ATEAKQK VDA**    153-168  
    **AELLENQ VHRLEQE LKEIDES** 181-  
 198  
    +  
                          **CNleG-IXXLXXX IXXLXXX IXXLXXX IXXLXXX IXXLXXX**  
    coiled-coil template  
 10     —  
                  **CNleG IEELEKK ITELKQK I--LENQ IHRLEQE IKELDES**  
    —                    fill gap with aa from either sequence  
                  **CNleG IEELEKK ITELKQK IDALENQ IHRLEQE IKELDES**  
 15     (test peptide; SEQ ID NO:3)

The peptide is synthesized as described in Materials and Methods. Also synthesized is a control peptide containing the native sequence instead of isoleucine and leucine at the a and d positions:

20                   **CNleG LEEAEKK ATEAKQK VDALENQ VHRLEQE LKEIDES**  
    (native control; SEQ ID NO: 4)

25                   The CD profile and helical stability of the peptides are determined as described in Materials and Methods. The results show that the test peptide exists in helical form even in benign buffer (50 mM KH<sub>2</sub>PO<sub>4</sub>, 50 mM KCL, pH 7.0) in the absence of trifluoroethanol (TFE). By contrast, the native control needs the presence of TFE to form a helical structure. Similarly, the test peptide is much

-42-

more stable than the native control when tested in a thermal denaturation experiment.

## Example 2 A synthetic pneumococcal PspA peptide

5

A synthetic peptide was prepared based on epitopes from three different Pneumococcal surface proteins A (PspA). The three PspA proteins and their capsular serotype are as follows:

	<u>PspA Strain</u>	<u>Capsular Serotype</u>
10	BG 8743	23F
	EF5668	4
	BG8090	19F

These proteins were chosen because of the prevalence of these serotypes in pneumococcal infections (Kalin, 1998). B-cell epitopes, shown in Figure 4, were identified using the bioinformatics software Peptools (Edmonton, Alberta). The three epitopes were spliced together such that the coiled-coil repeat remained in register. A linker and cysteine residue was incorporated to enable the strands of the coiled-coil to be linked together by a disulfide bond.

20

The CD spectrum of the peptide is shown in Figure 5. In benign buffer (50 mM KH<sub>2</sub>PO<sub>4</sub>, 50 mM KCl, pH 7.0) the peptide adopts a conformation that is a mix of random coil and alpha helix. In the presence of 50% trifluoroethanol (TFE, which induces helical structure in peptides that have the propensity to fold into an alpha helix), the peptide adopts a highly helical conformation. The helical content in benign buffer is 17% of that in 50% TFE, indicating that the peptide forms helices in a physiologically compatible solution without TFE. Therefore,

25

-43-

the peptide may be used to elicit an immune response under physiological conditions.

5     **Example 3     Generation of a consensus coiled-coil sequence**

10     To date 40 partial and full sequences of pneumococcal surface protein A from various strains have been released to various sequence databases. These proteins share a common architecture consisting of a leader sequence, a coiled-coil domain, a proline domain and a choline binding domain. Analysis of PspA sequences by the program MultiCoil (Wolf et al., 1997) indicates that the coiled-coil regions are broken up by sections which contain proline residues. Multiple sequence analysis using the software Biotools indicates that the published sequences vary in similarity from 24.1 % to 97 %. However, certain regions of the  
15     helical domain of these proteins are remarkably similar, suggesting that these conserved regions are important to the function of the protein

20     A consensus sequence (Figure 6) was deduced from a conserved region which was shown to have antigenic activities. Complete and partial PspA sequences were downloaded from the National Library of Medicine PubMed web site into the bioinformatics software Peptools (Biotools, Edmonton, Alberta). A total of 40 complete and partial sequences were transferred to the alignment module of the software and a consensus threshold was set to 65 %. The consensus threshold defines the minimum residue plurality amongst a group of aligned  
25     sequences. In effect, the consensus threshold acts to filter out insignificant matches and highlights conserved residues within a sequence. This allows for easy identification of similarities by visual inspection.



-44-

The last 100 amino acids of the coiled-coil region were the focus of sequence similarities since this region has been shown previously to harbor cross reactive epitopes. Figure 6 illustrates the sequence alignment for residues found within this region. In particular, the alignment yielded a consensus sequence of  
5 EELX<sub>1</sub>X<sub>2</sub>KIDELDX<sub>3</sub>EIAX<sub>4</sub>LEKX<sub>5</sub> (SEQ ID NO:5), in which the putative a and d positions were selected to be isoleucine and leucine, respectively. Preferably, X<sub>1</sub> is S, Q, N or D; X<sub>2</sub> is D, N or K; X<sub>3</sub> is A or N; X<sub>4</sub> is K, E or D; and X<sub>5</sub> is N, D or E.

10 A sequence containing the first section of this consensus region, EELSDKIDELD (SEQ ID NO:6), was selected for antigenicity studies. The Ser and Asp residues were included into this sequence since these residues appear in the majority of sequences aligned.

15 A synthetic peptide containing EELSDKIDELD (SEQ ID NO:6) was made according to formula I of the present invention. Thus, an isoleucine was inserted at the N-terminus of the above sequence according to formula I and flanking sequences were added to increase the length of the peptide. The complete sequence of this peptide, CVX0270, is as follows, wherein the residues from the  
20 consensus sequence are highlighted in bold script:

Acetyl-CNleGEIEALKKKIEEL**SDKIDELEKEIK**-amide (SEQ ID NO:7)

The CD spectra were performed under benign conditions (50 mM  
25 KH<sub>2</sub>P0<sub>4</sub>, 100 mM KCl, pH 7.0) and in aqueous buffer containing 50% TFE. The CD spectrum of peptide CVX0270 under benign conditions at 20 °C is shown in Figure 7. The CD spectra is typical of a helical peptide with minima at 222 nm

-45-

and 209 nm and high positive ellipticity below 200 nm. Typically, the molar ellipticity at 222 nm ( $[\theta]_{222}$ ) has been used to measure helical content in peptide. For peptide CVX0270 this corresponds to a value of -29800 which indicates that this peptide is predominantly  $\alpha$ -helical. Theoretically, the  $[\theta]_{222}$  value for a  
5 peptide of 27 residues is -33900 (Chen et al., 1974). Therefore peptide CVX0270 is 88% helical. It should be noted that the linker region (Cys-Nle-Gly) is designed not to be helical and therefore decreases the  $[\theta]_{222}$  signal. The ratio of the molar ellipticity at 222 and 208 ( $[\theta]_{222}/[\theta]_{208}$ ) is greater than 1.02 and similar to that observed before for coiled-coils (Hodges et al., 1988; Lau et al, 1984; Zhou et al.,  
10 1992) and distinctly different from non interacting  $\alpha$ -helices in which the  $[\theta]_{208}$  is greater than the  $[\theta]_{222}$ . In the presence of 50% TFE, helical content increased slightly to 101%. The data indicate that peptide CVX0270 is highly helical and found predominantly in the coiled-coil conformation under aqueous conditions.

15 In order to determine the stability of CVX0270, a thermal denaturation study was undertaken. Figure 8 shows the denaturation curve obtained by monitoring the  $[\theta]_{222}$  as a function of temperature. The study indicates that peptide CVX0270 is very stable with the peptide exhibiting 77% of its original helicity at 75 °C. The above results demonstrate that the coiled-coil forming sequence with  
20 isoleucine at a positions and leucine residues at d positions is sufficiently stable to house a helical epitope from another protein sequence.

25

-46-

**Example 4      Immunogenicity of Coiled-Coil Proteins**

The peptide "Strep" (SEQ ID NO:7) as described in Example 3 was coupled to the protein carrier tetanus toxoid (TT) as described in Materials and  
5 Methods. Immunogenicity of this peptide was determined as follows.

Balb/c mice (6-8 weeks, female, Charles River) were immunized on day 0 (1° primary immunization), day 7 (2° secondary immunization), day 28 (3° tertiary immunization), and day 42 (boost) by intraperitoneal injection ( 100  $\mu$ L total, 50%  
10 Alhydrogel 2% adjuvant, Cedarlane, Catalog # SF2000-250) with Strep-tetanus toxoid (20, 10 or 5  $\mu$ g) and with unconjugated tetanus toxoid. These antigens were diluted to various doses in 0.9% NaCl and mice injected with 0.9% NaCl were used as negative controls. Mice were bled following the 1°, 2°, 3° and boost injections on days 6, 14, 35 and 49 to collect serum to assay for antibody titers  
15 (direct ELISA) and antibody isotype response (isotyping ELISA), determine antibody specificity (inhibition ELISA) and measure immunoprotective antibody responses (bactericidal/opsonization assays).

A typical immunization schedule is shown in Table 1. Various other  
20 immunization schedules and adjuvant formulations known in the art would also be effective. Various delivery methods including intravascular, intramuscular, intradermal, subcutaneous, oral, nasal and aerosol inhalation routes would also be effective.

25

**Table 1: BALB/c Mouse immunized I.P. – Strep-TT**  
**Immunization Schedule**

5	<b>Day</b>	<b>Procedure</b>
	0	Immunize i.p.
10	6	Bleed
	7	Immunize i.p.
	14	Bleed
	28	Immunize i.p.
	35	Bleed
	42	Immunize i.p.
	49	Cardiac Bleed

15

**Study Groups:**  
**(10 A.BY mice / group)**

1. Strep-TT (20  $\mu$ g/mouse) + adjuvant
202. Strep- TT ( 10  $\mu$ g / mouse) + adjuvant
3. Strep- TT (5  $\mu$ g / mouse) + adjuvant
254. Tetanus toxoid (20  $\mu$ g/mouse) + adjuvant
5. PBS (without Alum)

30

-48-

A. Determining Antibody Levels Elicited by Coiled-Coil Antigen Conjugates

5 The basic procedure to measure antibody levels is by the following direct ELISA protocol:

1. Coat EIA plates (COSTAR) with 1.0  $\mu\text{g/mL}$  (0.1  $\mu\text{g}$  per well) of antigen in 0.05 M carbonate-bicarbonate buffer (pH 9.6, 100  $\mu\text{L}$ /well).
- 10 2. Incubate the plate(s) at 4°C. overnight.
3. On the next day, wash plates 3X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.
- 15 4. Block plates with 100  $\mu\text{L}$  blocking buffer per well (PBS / 2% BSA). Incubate plates for 1 hour at 37°C.
5. Wash plates 5X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.
- 20 6. Add 100  $\mu\text{L}$  per well of test antibody appropriately diluted in dilution buffer (PBS / 0.1% Tween). Incubate plates for 60 minutes at 37°C.
7. Wash plates 3X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.
- 25

-49-

8. Dilute HRP anti-mouse IgG (Jackson Lab) in dilution buffer (PBS / 0.1 % Tween) to a concentration of 1:5000. Add 100  $\mu$ L per well and incubate at 37°C for 60 minutes.
- 5 9. Wash plates 3X with washing buffer (PBS / 0.05 % Tween). Flick off excess liquid by tapping the plates on the bench top.
- 10 10. Prepare HRP substrate;  
0.03 % Hydrogen peroxide (30%) and 1 mM ABTS dissolved in 0.01 M Sodium citrate buffer (Sodium citrate (1.48 g) dissolve sodium citrate in 500 mL distilled H<sub>2</sub>O. Citric acid (1.58 g) dissolve citric acid in 750 mL distilled H<sub>2</sub>O. Pour citric acid solution into sodium citrate solution to get a pH 4.2). Add 100  $\mu$ L to each well and develop in a dark place for 30 minutes.
- 15 11. Read the absorbance with an ELISA plate reader at 405 nm at 30 minutes.

The mean O.D. ELISA readings obtained with antisera to the Strep-TT conjugates (20  $\mu$ g, 10  $\mu$ g and 5  $\mu$ g) is shown in Table 2. The Strep-TT conjugate at all three dose concentrations (20  $\mu$ g, 10  $\mu$ g and 5  $\mu$ g) elicited significant antibody titers to the Strep (EIEALKKKIEELSDKIDELEKEIK) hapten. These antibodies were immunogen specific, as the unconjugated tetanus toxoid did not elicit any antibodies or cross-reactive antibodies to the coiled-coil Strep antigen.

25



-50-

**Table 2: ELISA Results of Mouse Serum Antibodies to the  
Coiled-Coil Strep Antigen**

5	Serum	Murine Sera (Day 49) (OD 405 nm, Strep-BSA coating antigen)				
	Dilutions	Strep-TT	Strep-TT	Strep-TT	TT*	PBS*
		(20 $\mu$ g)*	(10 $\mu$ g)*	(5 $\mu$ g)*		
10	<b>1:2000</b>	3.75	3.71	3.86	0.11	0.01
	<b>1:4000</b>	3.57	3.60	3.61	0.04	0.00
	<b>1:8000</b>	2.48	2.66	2.60	0.01	0.00
	<b>1:16000</b>	1.14	1.59	1.50	0.06	0.00
	<b>1:32000</b>	0.76	0.84	0.95	0.00	0.00
	<b>1:64000</b>	0.36	0.42	0.41	0.00	0.00
	<b>1:128000</b>	0.18	0.23	0.22	0.00	0.00
15	<b>1:256000</b>	0.04	0.10	0.12	0.00	0.00

\* mice were i.p. injected on day 0, 7, 28 and 42 with Strep-TT +  
adjuvant, TT + adjuvant or PBS. Sera were collected on day 49.

-51-

B. Determining Antibody Isotype Levels Elicited by Coiled-Coil Antigen Conjugates

5           The basic procedure to measure antibody isotype levels is as follows to quantify IgM, IgG, and IgA isotypes elicited by coil-coiled antigen conjugates.

1. Coat EIA plates (COSTAR) with 1.0  $\mu\text{g/mL}$  (0.1  $\mu\text{g}$  per well) of antigen in 0.05 M carbonate-bicarbonate buffer pH 9.6, 100  $\mu\text{L}$ /well.

10

2. Incubate the plate(s) at 4°C overnight.

3. On the next day, wash plates 3X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.

15

4. Block plates with 100  $\mu\text{L}$  blocking buffer per well (PBS / 2% BSA). Incubate plates for 1 hour at 37°C.

20

5. Wash plates 5X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.

6. Prepare 1:250 dilution mouse serum in the working buffer (PBS / 0.1% Tween). Add 100  $\mu\text{L}$ /well into the appropriate well. Incubate plates for 1 hour at 37°C.

25

7. Wash plates 3X with washing buffer (PBS / 0.05% Tween). Flick off excess liquid by tapping the plates on the bench top.

-52-

8. Dilute HRP-labeled detection antibodies (Southern Biotechnology Associates Inc.). add 100  $\mu$ L/well into the appropriate well. Incubate plates for 1 hour at 37°C.
- 5
9. Wash plates 3X with washing buffer (PBS / 0.05 % Tween). Flick off excess liquid by tapping the plates on the bench top.
10. Prepare HRP substrate;
- 10      0.03 % Hydrogen peroxide (30 %) and 1mM ABTS dissolved in 0.01 M Sodium citrate buffer (Sodium citrate (1.48 g) dissolve sodium citrate in 500 mL distilled H<sub>2</sub>O. Citric acid (1.58 g) dissolve citric acid in 750 mL distilled H<sub>2</sub>O. Pour citric acid solution into sodium citrate solution to get a pH 4.2). Add 100  $\mu$ L to each well and develop in a dark place for 30 minutes
- 15
11. Read the absorbance with an ELISA plate reader at 405 nm at 30 minutes.

The IgG antibody isotype response to the Strep-TT coiled-coil antigen is shown in Table 3. These results demonstrate the presence of significant IgG1 antibody levels to the Strep coiled-coil antigen after secondary immunization. A significant IgG isotype switch from IgG1 to IgG<sub>2a</sub>, IgG<sub>2b</sub> and IgG<sub>3</sub> was seen after the booster injection (Day 49 bleed). The observed IgG antibody maturation to the coiled-coil Strep antigen is typical of a TD response.

20

-53-

**Table 3: Isotyping ELISA of Mouse Serum Antibodies to the Coiled-Coil Strep Antigen**

**DAY 6**

Antiserum to:*	Isotypes (O.D. 405 nm)			
	IgG <sub>1</sub>	IgG <sub>2a</sub>	IgG <sub>2b</sub>	IgG <sub>3</sub>
Strep-TT (20 $\mu$ g)	0.14	0.01	0.04	0.04
Strep-TT (10 $\mu$ g)	0.12	0.01	0.03	0.04
Strep-TT (5 $\mu$ g)	0.14	0.05	0.08	0.10
TT (20 $\mu$ g)	0.06	0.02	0.01	0.01
PBS	0.03	0.00	0.00	0.01

\* mice were i.p. injected on day 0 with Strep-TT + adjuvant, TT + adjuvant or PBS. Sera were collected on day 6.

**DAY 14**

Antiserum to:*	Isotypes (O.D. 405 nm)			
	IgG <sub>1</sub>	IgG <sub>2a</sub>	IgG <sub>2b</sub>	IgG <sub>3</sub>
Strep-TT (20 $\mu$ g)	2.00	0.71	0.95	1.08
Strep-TT (10 $\mu$ g)	1.84	0.91	1.01	2.22
Strep-TT (5 $\mu$ g)	1.69	0.62	0.95	1.23
TT (20 $\mu$ g)	0.05	0.00	0.00	0.01
PBS	0.03	0.00	0.00	0.00

\* mice were i.p. injected on day 0 and 7 with Strep-TT + adjuvant, TT + adjuvant or PBS. Sera were collected on day 14

-54-

**Day 35**

	Antiserum to:*	Isotypes (O.D. 405 nm)			
		IgG <sub>1</sub>	IgG <sub>2a</sub>	IgG <sub>2b</sub>	IgG <sub>3</sub>
5	Strep-TT (20 µg)	2.24	0.40	0.49	0.45
	Strep-TT (10 µg)	2.07	0.56	0.59	1.00
	Strep-TT (5 µg)	2.16	0.60	0.76	0.75
	TT (20 µg)	0.08	0.01	0.01	0.01
	PBS	0.01	0.00	0.00	0.01

10 \* mice were i.p. injected on day 0, 7 and 28 with Strep-TT +  
adjuvant, TT + adjuvant or PBS. Sera were collected on day 35.

**Day 49**

	Antiserum to:*	Isotypes (O.D. 405 nm)			
		IgG <sub>1</sub>	IgG <sub>2a</sub>	IgG <sub>2b</sub>	IgG <sub>3</sub>
15	Strep-TT (20 µg)	3.91	1.14	1.28	0.74
	Strep-TT (10 µg)	3.85	1.35	2.01	1.28
	Strep-TT (5 µg)	3.88	2.10	2.96	0.82
20	TT (20 µg)	0.44	0.03	0.02	0.05
	PBS	0.01	0.01	0.01	0.02

\* mice were i.p. injected on day 0, 7, 28 and 42 with Strep-TT +  
adjuvant, TT + adjuvant or PBS. Sera were collected on day 49.

25

-55-

C. Determining Antibody Binding in Competition with Coiled-Coil Antigen Conjugates or Free Peptides

5           The specificity of the antisera was further examined by inhibition ELISA using peptides with or without a coiled-coil structure. The protocol for this inhibition ELISA is as follows:

10           1. Dilute the coating antigen to 1.0  $\mu\text{g/mL}$  (0.1  $\mu\text{g}$  per well) in carbonate-bicarbonate buffer. Use glass tubes.

2           Add 100  $\mu\text{L}$  of the coating antigen to each well of the plate. Store the plate(s) overnight at 4°C.

15           3. Shake out the wells and tap upside-down onto Kimwipes. Wash plate three times with  $\sim 200 \mu\text{L}$  wash buffer (PBS / 0.05% Tween)(v/v) per well.

20           4 Add 200  $\mu\text{L}$  blocking buffer per well (PBS / 2% BSA)(w/v). Incubate plates for 60 minutes at 37°C.

25           5. Shake out the wells and tap upside-down onto Kimwipes. Wash plate five times with  $\sim 200 \mu\text{L}$  wash buffer (PBS / 0.05% Tween)(v/v) per well.



-56-

6. Add 50  $\mu\text{L}$  of selected competitive conjugates or free peptides appropriately diluted in dilution buffer (PBS / 0.1 % Tween)(v/v). (NOTE: the competitive peptide has a x2 concentration in 50  $\mu\text{L}$ ).
- 5 7. Added 50  $\mu\text{L}$  per well of test antibody appropriately diluted in dilution buffer (PBS / 0.1 % Tween)(v/v). (NOTE: the test antibody has a x2 concentration in 50  $\mu\text{L}$ ). The final volume in each well was 100  $\mu\text{L}$ .
- 10 8. Incubate plates for 60 minutes at 37°C with slow shaking on a ELISA shaker.
9. Shake out the wells and tap upside-down onto Kimwipes. Wash plate three times with  $\sim 200$   $\mu\text{L}$  wash buffer (PBS / 0.05 % Tween)(v/v) per well. Dilute Peroxidase-conjugated anti-mouse IgG  
15 in dilution buffer (PBS / 0.1 % Tween)(v/v) to a concentration of 1:5000. Add 100  $\mu\text{L}$  per well and incubate at 37°C for 60 minutes.
10. Shake out the wells and tap upside-down onto Kimwipes.  
20 Wash plate three times with  $\sim 200\mu\text{L}$  wash buffer (PBS / 0.05 % Tween)(v/v) per well.
11. Prepare HRP substrate;  
0.03 % Hydrogen peroxide (30%) and 1mM ABTS dissolved in  
25 0.01M Sodium citrate buffer (Sodium citrate (1.48 g) dissolve sodium citrate in 500 mL distilled  $\text{H}_2\text{O}$ . Citric acid (1.58 g) dissolve citric acid in 750 mL distilled  $\text{H}_2\text{O}$ . Pour citric acid solution into sodium

-57-

citrate solution to get a pH 4.2).. Add 100  $\mu$ L to each well and develop in a dark place for 30 minutes

12. Read the absorbance with an ELISA plate reader at 405 nm at  
5 30 minutes.

The sequences of the Strep #1, #2 and #3 free peptides used in this inhibition assay are:

10 Strep #1: Coiled-coil Strep antigen with lactams  
**EIEALKKKIEELSDKIDELEKEIK**

Strep #2: Single stranded Strep antigen with lactams  
**EEESDKKDEEDAEKAE**

15 Strep #3: Coiled coil Strep antigen  
**EIEALKKKIEELSDKIDELEKEIK**

20 The residues that make up the lactam bridges are highlighted in a bold font. The lactams are formed by reacting the carboxylic acid side chains of glutamic acids residues with the side chain amino groups of lysine residues. These lactams stabilize  $\alpha$ -helices and coiled coils when they are situated 4 residues apart and when they are oriented in the glutamic acid to lysine direction.

25

The epitope is highlighted in italicized script. From all three sequences it can be seen that the epitope begins with two glutamic acid residues (**EE**)

-58-

followed by a leucine residue (L). In the case of the single stranded peptide the leucine residue is replaced by a glutamic acid (E) residue which is part of the lactam bridge. The epitope continues with **SKD** followed by an isoleucine (I) in the coiled-coils and the lysine residue (K) which is involved in the lactam bridge in the single stranded peptide. The final two residues of the epitope are **DE**.

Table 4 shows the inhibition ELISA results of anti-Strep serum using 3 different Strep peptides, the Strep-TT immunogen, TT and BSA. The Strep #3 peptide (peptide used as the immunogen) showed significant inhibition, the Strep #2 peptide (linear peptide, which has low helical content) showed no inhibition. Therefore, it would seem that the immunogen Strep-TT elicited antibodies which recognize conformational (or 3D) epitope structures. The Strep-TT immunogen did not elicit antibodies to the linear epitope. These results therefore suggest that coiled-coil antigens elicit antibodies which recognize conformational epitopes. Inhibition was also seen with Strep #1 peptide, which peptide contains lactam bridges which adds further stability to the coiled-coil antigen.

-59-

**Table 4: Inhibition ELISA of Anti-Strep.(1:25,000 dilution, Day  
49) Using Various Strep Antigen Inhibitors**

5	<b>Anti-Strep.-T.T. (20<math>\mu</math>g Dose)</b>						
	O.D. 405 nm						
10	<b>Conc.</b> ( $\mu$ g per 100 $\mu$ L)	<b>Strep.</b> <b>#1</b>	<b>Strep.</b> <b>#2</b>	<b>Strep.</b> <b>#3</b>	<b>Strep.-</b> <b>T.T.</b>	<b>T.T.</b>	<b>BSA</b>
	<b>10</b>	0.663	1.266	0.110	0.028	1.342	1.399
15	<b>5</b>	0.720	1.186	0.143	0.026	1.051	1.485
	<b>2.5</b>	0.823	1.173	0.254	0.056	1.412	1.435
20	<b>1</b>	0.959	1.164	0.555	0.094	1.457	1.442
	<b>0.5</b>	1.081	1.144	0.776	0.109	1.564	1.187
25	<b>0.25</b>	1.237	1.251	1.036	0.165	1.638	1.510
30	<b>Anti-Strep.-T.T. (10<math>\mu</math>g Dose)</b>						
	O.D. 405 nm						
35	<b>Conc.</b> ( $\mu$ g per 100 $\mu$ L)	<b>Strep.</b> <b>#1</b>	<b>Strep.</b> <b>#2</b>	<b>Strep.</b> <b>#3</b>	<b>Strep.-</b> <b>T.T.</b>	<b>T.T.</b>	<b>BSA</b>
	<b>10</b>	0.941	1.210	0.055	0.010	1.820	1.784
40	<b>5</b>	0.917	1.258	0.233	0.010	1.521	1.790
	<b>2.5</b>	0.963	1.232	0.269	0.015	1.726	1.765
45	<b>1</b>	1.129	1.312	0.619	0.065	1.802	1.670
	<b>0.5</b>	1.256	1.293	0.872	0.068	1.800	1.422
50	<b>0.25</b>	1.340	1.279	1.123	0.113	1.871	1.845

-60-

	Anti-Strep.-T.T. (5 $\mu$ g Dose)						
	O.D. 405 nm						
	Conc. ( $\mu$ g per 100 $\mu$ L)	Strep. #1	Strep. #2	Strep. #3	Strep.- T.T.	T.T.	BSA
5	10	0.872	1.294	0.009	0.000	1.803	1.740
10	5	0.960	1.318	0.090	0.000	1.109	1.571
	2.5	1.087	1.233	0.213	0.000	1.822	1.761
	1	1.125	1.322	0.501	0.053	1.797	1.572
	0.5	1.289	1.377	0.746	0.064	1.836	1.717
	0.25	1.388	1.303	1.001	0.107	1.925	1.786

15

Strep. #1 = Coiled-coil + lactam bridges

Strep. #2 = Linear sequence

Strep. #3 = SEQ ID NO:7

20

-61-

D. Bactericidal and Opsonization Assays to Measure Immunoprotective Antibodies Elicited by Coil-Coiled Antigen Conjugates

5           The basic bactericidal and opsonization assays used are as follows:

**Bactericidal Assay**

1. Streak an agar plate with desired gram positive bacteria procured from various culture collections (ATCC). Incubate at 37°C overnight.
- 10           2. On the next day, pick an isolated colony and inoculate it in 1.0 ml of Todd-Hewitt Broth (THB) + Yeast Extraction (YE) media in a sterile test tube. Incubate at 37°C overnight.
- 15           3. On the following day, measure O.D. of inoculated bacteria at 420 nm wavelength. Use THB + YE media as blank.
4. To a sterile flat bottom 96-well plate, add a sterile 2.5 mm glass bead in each well.
- 20           5. To each well, add:  
            5  $\mu$ L of bacteria  
            10  $\mu$ L of mouse serum to be tested  
            incubate at 37°C for 1 hour.
- 25           NOTE: Step 5 and 6 are done in triplicate.



-62-

6. After 1 hour incubation, prepare 1:20 dilution exogenous complement (Low-Tox guinea pig Complement, Cedarlane) sterilely in THB + YE. Add 50  $\mu$ L/well. Incubate at 37°C for 1 hour.
- 5     7. After complement incubation, 50  $\mu$ L aliquot is plated out on agar plates using glass spreader.
8. Wrap all agar plates in plastic bags and incubate at 37°C. overnight.
- 10    9. On the next day, count colony forming units (CFU).

#### **Opsonization Assay**

- 15     1. Streak an agar plate with desired gram positive bacteria procured from various culture collections (ATCC). Incubate at 37°C overnight.
2. On the next day, pick an isolated colony and inoculate it in 1.0 ml of Todd-Hewitt Broth (THB) + Yeast Extraction (YE) media in a sterile test  
20     tube. Incubate at 37°C overnight.
3. The next day prepare 100 U/ml of sterile heparin.
4. I.V. inject 100  $\mu$ L of sterile heparin into tail vein of each mouse (5-10).  
25     After 10 minutes, cardiac bleed mice into a sterile tube.

-63-

5. Measure O.D. of inoculated bacteria at 420 nm wavelength. Use THB + YE media as blank.

5 6. To a sterile flat bottom 96-well plate, add a sterile 2.5 mm glass bead in each well.

7. To each well, add:

- a 50  $\mu$ L of heparinized blood
- b 10  $\mu$ L of mouse serum to be tested
- c 5  $\mu$ L of bacteria

incubate at 37°C for 1 hour.

NOTE: Do this step in triplicate.

15 8. After 1 hour incubation, prepare 1:20 dilution exogenous complement (Low-Tox guinea pig Complement, Cedarlane) sterilely in distilled water. Add 50  $\mu$ L/well. Incubate at 37°C for 1 hour on a shaker (slow motion).

9. After one hour, 100  $\mu$ L aliquot is plated out on agar plates using glass spreader.

20

10. Wrap all agar plates in plastic bags and incubate at 37°C overnight.

11. On the next day, count colony forming units (CFU).

25

Sera from mice immunized with the Strep-TT conjugates (20, 10, 5  $\mu$ g doses) were found to be immunoprotective as measured by the opsonization assay.

Antisera to Strep-TT showed reduction in colony forming units of representative *Streptococcus pneumoniae* serotypes (1, 3, and 5). Little or no reduction in colony forming units was seen with sera from mice injected with tetanus toxoid or with PBS. The Strep-TT conjugate elicited immunoprotective antibodies to coiled-coil antigens of *S. pneumoniae*.

**Table 5: Opsonization Results of Antiserum to Strep-TT (Day 49) using *Streptococcus pneumoniae* serotypes (1, 3 and 5)**

Percent Reduction of CFU			
Antiserum to:	Type 1	Type 3	Type 5
Strep-TT (20 µg)	58	49	45
Strep-TT (10 µg)	22	50	24
Strep-TT (5 µg)	30	48	24
TT	7	6	7
PBS	0	0	1

-65-

## Appendix 1

5

Analysis of immunogenic regions of PspA proteins from *S. pneumoniae* strain Rx1. Immunogenic regions are indicated by asterisks underneath the sequence.

-66-

Name: PspA Rx1

Organism: S. pneumoniae

	10	20	30	40	50	
1	EESPVASQSKAEKDYDAAKKDAKNAKKAVEDAQKALDDAKAAQKKYDEDQ					50
	5*****					52
	3*****					29*****
	10	20	30	40	50	
51	KKTEEKAALKAASEEMDKAVAAVQQAYLAYQQATDKAAKDAADKMIDEA					100
	***** 66***** 73*****					
						78**
	10	20	30	40	50	
101	KKREEEAKTKFNTVRAMVVPEPEQLAETKKKSEEAKQKAPELTKKLEEAK					150
	91***** 95***** 123*****					
	**** 101**** 122*****					
	89***** 116*****					
	10	20	30	40	50	
151	AKLEEAEEKKATEAKQKVDAEEVAPQAKIAELENQVHRLEQELKEIDSESES					200
	133***** 140***** 148***					
	130***** 145*****					
	10	20	30	40	50	
201	EDYAKEGFRAPLQSKLDAKKAKLSKLEELSDKIDELDAEIAKLEDQLKAA					250
	* 168*****					
	166***** 1					
	10	20	30	40	50	
251	EENNNVEDYFKEGLEKTIAAKKAEELEKTEADLKKAVNEPEKPAPAPETPA					300
	179***** 185***** 195***** 206*****					
	76***** 182*****					
	10	20	30	40	50	
301	PEAPAEQPKPAPAPQPAPAPKPEKPAPAEQPKPEKTDDQQAEEEDYARRSEEE					350

-67-

	10	20	30	40	50	
351	YNRLTQQQPPKAEKPAPAPKTGWKQENGMWYFYNTDGSMATGWLQNNGSW					400
401	YYLNSNGAMATGWLQYNGSWYYLNANGAMATGWAKVNGSWYYLNANGAMA					450
451	TGWLQYNGSWYYLNANGAMATGWAKVNGSWYYLNANGAMATGWLQYNGSW					500
501	YYLNANGAMATGWAKVNGSWYYLNANGAMATGWVKDGD TWYYLEASGAMK					550
551	ASQWFKVSDKWYYVNLGLALAVNTTVDGYKVNANGEWV					588

---

Motif #: 3 Range: 6 - 13  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 5 Range: 8 - 15  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 29 Range: 42 - 49  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST]

-68-

ST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 52 Range: 49 - 56

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 66 Range: 63 - 70

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 73 Range: 81 - 88

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 78 Range: 97 - 104

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 89 Range: 104 - 111

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 91 Range: 107 - 114

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN



-69-

Motif #: 95 Range: 119 - 125  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 101 Range: 127 - 133  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 116 Range: 132 - 139  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 122 Range: 139 - 146  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 123 Range: 142 - 149  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 130 Range: 153 - 160  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED  
Name: GENERALIZED B CELL EPITOPE PATTERN

---

Motif #: 133 Range: 160 - 167  
Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*  
Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 140 Range: 181 - 188

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 145 Range: 192 - 199

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 148 Range: 195 - 201

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 166 Range: 226 - 233

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 168 Range: 229 - 236

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 176 Range: 250 - 256

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 179 Range: 252 - 259

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

-71-

ST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 182 Range: 260 - 267

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 185 Range: 261 - 268

Motif: \*[DEGHKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 195 Range: 275 - 282

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

Motif #: 206 Range: 288 - 295

Motif: \*[DEGHKNPQRST][DEHGKNPQRST][DEHGKNPQRST][DEHGKNPQRST]\*[DEHGKNPQRST]\*

Ref.: WISHART, D.S. UNPUBLISHED

Name: GENERALIZED B CELL EPITOPE PATTERN

-72-

We claim:

1. A synthetic peptide of the formula I:



wherein

- 5        A is Ile, Leu, Val or a derivative thereof;  
      D is Leu, Ile, Val or a derivative thereof;  
          each X is an amino acid residue or derivative thereof which  
          corresponds to an amino acid residue of an epitope of a native  
          coiled-coil protein;  
10        the X residues in each (AXXDXXX) repeat form a set of X residues; and  
          n is equal to or greater than 1.
- 15        2. The peptide of Claim 1 wherein A is Ile and D is Leu in every  
          (AXXDXXX) repeat.
3. The peptide of Claim 1 wherein n is about 3 to 6.
4. The peptide of Claim 1 wherein said X residues are amino acids that are  
          solvent exposed in an coiled-coil region of the native protein.
- 20        5. The peptide of Claim 1 wherein each of said sets of X residues is from the  
          same epitope of a single protein.
6. The peptide of Claim 1 which contains at least two different sets of X  
25        residues.

-73-

7. The peptide of Claim 6 wherein each of said different sets is independently selected from the group consisting of different epitopes of the same protein and epitopes from different proteins.
- 5 8. The peptide of Claim 1 which further comprises additional amino acids at the C-terminus and/or N-terminus of the peptide.
9. The peptide of Claim 8 wherein said additional amino acid residues are CNleG at the N-terminus of the peptide.
- 10 10. The peptide of Claim 1 wherein the set of X residues correspond to a consensus sequence of solvent exposed residues of native coiled-coil proteins.
- 15 11. The peptide of Claim 10 wherein the coiled-coil proteins are selected from the group consisting of Pneumococcal surface protein A, Pneumococcal surface protein C, and Pneumococcal adhesin A.
- 20 12. The peptide of Claim 11 wherein the peptide comprises an amino acid sequence selected from the group consisting of EELX<sub>1</sub>X<sub>2</sub>KIDELDX<sub>3</sub>EIAX<sub>4</sub>LEKX<sub>5</sub> (SEQ ID NO:5) and EELX<sub>1</sub>X<sub>2</sub>KIDELD (1-11 of SEQ ID NO:5), wherein X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub>, X<sub>4</sub> or X<sub>5</sub> is any amino acid.
- 25 13. The peptide of Claim 12 wherein  
X<sub>1</sub> is S, Q, N or D;  
X<sub>2</sub> is D, N or K;  
X<sub>3</sub> is A or N;

-74-

X<sub>4</sub> is K, E or D; and  
X<sub>5</sub> is N, D or E.

14. A synthetic peptide of the formula I:

5 (AXXDXXX)<sub>n</sub> I

wherein

A is Ile, Leu, Val or a derivative thereof;

D is Leu, Ile, Val or a derivative thereof;

10 each X is an amino acid residue or derivative thereof which  
corresponds to an amino acid residue of an epitope of a native  
coiled-coil protein, except at least one X is replaced with a charged  
amino acid residue in a manner which allows a salt bridge to form  
between the charged amino acid and another amino acid residue of  
an opposite charge, which salt bridge facilitates the peptide to  
15 assume a coiled-coil structure;

the X residues in each (AXXDXXX) repeat form a set of X residues; and  
n is equal to or greater than 1.

15. A peptide of Claim 14 wherein the charged amino acid is selected from the  
20 group consisting of Asp, Glu, Lys, Arg and His.

16. A method of making a peptide of the formula I comprising:

- 25 a) selecting an epitope of a coiled-coil protein;  
b) determining which amino acid residues of said epitope are solvent exposed;  
and  
c) inserting said solvent exposed amino acid residues into the X positions of  
formula I.

-75-

17. The method of Claim 16 wherein the coiled-coil protein is a microbial protein.
- 5 18. The method of Claim 16 wherein the selection of epitopes is performed using a computer algorithm.
19. The method of Claim 16 wherein more than one set of epitopic amino acids is used.
- 10 20. The method of Claim 19 wherein each of said sets is independently selected from the group consisting of different epitopes of the same protein and epitopes from different proteins.
- 15 21. A composition useful to stimulate an immune response in an animal, said composition comprising at least one peptide of formula I.
22. The composition of Claim 21 wherein the peptide of formula I is conjugated to a carrier protein.
- 20 23. The composition of Claim 21 further comprising an adjuvant.
24. The composition of Claim 21 which contains at least two different sets of X residues.
- 25



-76-

25. The composition of Claim 24 wherein each of said different sets is independently selected from the group consisting of different epitopes of the same protein and epitopes from different proteins.
- 5 26. The composition of Claim 24 which is useful to stimulate an immune response to more than one strain and/or species of microorganism.
27. A method of eliciting an immune response in an animal, comprising administering a peptide of the formula I to said animal.
- 10 28. An antibody which recognizes a peptide of the formula I.
29. The antibody of Claim 28 wherein the peptide of the formula I contains solvent exposed amino acids from a microbial protein.
- 15 30. The antibody of Claim 28 which binds to more than one strain and/or species of microorganism.
31. The antibody of Claim 28 which is polyclonal or monoclonal.
- 20 32. A pharmaceutical composition comprising an antibody according to Claim 28.
- 25 33. The composition of Claim 32 which further comprises a pharmaceutically acceptable excipient or carrier.

-77-

34. An antibody produced by administering a peptide of the formula I to an animal so as to stimulate an immune response.
- 5 35. A composition useful as a vaccine, wherein said composition comprises a peptide of formula I.
36. The composition of Claim 35 wherein more than one set of epitopic amino acids is used in the peptide of formula I.
- 10 37. The composition of Claim 36 wherein the sets of epitopic amino acids are from different strains and/or species of microorganism.
38. The composition of Claim 35 which provides cross protection to more than one strain and/or species of microorganism.
- 15 39. The composition of Claim 36 which provides cross protection to more than one strain and/or species of microorganism.
40. The composition of Claim 37 which provides cross protection to more than one strain and/or species of microorganism.
- 20 41. The composition of Claim 35 which further comprises a pharmaceutically acceptable excipient or carrier.
- 25 42. A method of preventing a microbial infection comprising administering to a mammal susceptible to said infection a peptide of formula I.

-78-

43. The method of Claim 42 wherein more than one set of epitopic amino acids is used in the peptide of formula I and the sets of epitopic amino acids are from different strains and/or species of microorganism.
- 5 44. The method of Claim 42 which is useful to prevent infection by several strains and/or species of microorganism.
45. The method of Claim 43 which is useful to prevent infection by several strains and/or species of microorganism.
- 10 46. A method of treating or preventing microbial infection in an animal susceptible to or suffering from such infection, comprising administering to said animal an effective amount of an antibody to a microbial protein, wherein said antibody is produced by administering a peptide of formula I
- 15 to an animal.
47. The method of Claim 46 which prevents symptoms of infection in said animal.
- 20 48. The method of Claim 46 which is useful to treat or prevent infection by several strains and/or species of microorganism.
49. A method of determining the presence of a particular microorganism in a sample comprising:
- 25 a) contacting the sample with an antibody to a peptide of formula I which peptide comprises epitopes from the particular microorganism; and

-79-

b) determining whether said antibody binds to a component of said sample.

50. The method of Claim 49 wherein the sample is a biological sample.

5

51. The method of Claim 49 which is used to determine the causative agent of a microbial infection.

10

52. The method of Claim 51 which is used to simultaneously detect the presence of several strains and/or species of microorganism in the sample.

53. The method of Claim 50 which is used to simultaneously detect the presence of several strains and/or species of microorganism in the sample.

15

54. The method of Claim 51 which is used to simultaneously detect the presence of several strains and/or species of microorganism in the sample.

55. A method for determining the presence of antibodies to a microbial protein in a biological sample, comprising:

20

a) contacting said biological sample with a peptide of formula I, which peptide comprises at least one epitope from said microbial protein; and

b) determining whether antibodies in said biological sample bind to said peptide.

25

56. The method of Claim 55 which is used to determine prior exposure of an animal to a particular microorganism.

-80-

Patent

Attorney Docket No. 003592-007

57. The protein of Claim 8 wherein the additional amino acids stabilize the peptide through the formation of lactam bridges.

5

10

1/8




Figure 1

2/8




Figure 2






Figure 3

4/8

1. Identify Potential epitopes in PspA from Clinically Revelant Strains

Strain BG 8743 (Sterotype 23F)  
 defgabcdefgabcdefgab  
 LEKDVEDFKNSDGEQAEQVL (SEQ ID NO: 10)  
 Residues 290-301

Strain EF5668 (Sterotype 4)  
 defgabcdefgabacd  
 LEDNLKDAETNNVED (SEQ ID NO: 11)  
 Residues 330-344

Strain BG 8090 (Sterotype 19F)  
 defgabcdef  
 LAKKQTELEK (SEQ ID NO: 12)  
 Residues 267-276

2. Splice Epitopes Together

DVEDFKNSDGELEDNLKDAETNNVEDAKKQTELEK (SEQ ID NO: 13)

3. Replace a and d Positions (underlined)

DIEDLKNSIGELEDNIKELETNIVELAKKITELEK (SEQ ID NO: 14)

**Figure 4**

5/8




Figure 5

6/8

Consensus	--E-----E-L--KIDELD-EIA-LEK-V---E-----K-----E---E-Y--	79
BG6692	--E-----E-LSDKIDELDAETAKLEKNV---E-DF-KNSNGEQ-AEOY-R	299
Strain 23	--E-----E-LSDKIDELDAETAKLEKNV---E-DF-KNSNGEQ-AEOY-R	135
Strain SP	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEY-SALY-L	135
Strain SP	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEY-SALY-L	135
BG8838	--E-----E-LSDKIDELDAETAKLEKNV---E-DF-KNSNGEQ-AEOY-R	299
strain 13	--E-----E-LSDKIDELDAETAKLEKNV---E-DF-KNSNGEQ-AEOY-R	95
Strain SP	--E-----A-LQNKVADLENKVAELDKEVTRLQSDL-KDAEENN-VEDYVK	127
Strain 18	--E-----A-LQNKVADLENKVAELDKEVTRLQSDL-KDAEENN-VEDYVK	98
BG8743	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	307
Strain 16	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	143
Strain 90	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	121
Strain 13	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	120
Strain 11	--A-----E-LDKKADELQNKVADLEKEISNLEILL-GGADPE---DDT--	86
Strain 43	--A-----E-LNKKVESLQNKVADLEKEISNLEILL-GGADPE---DDT--	99
Strain 23	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	84
BG11703	AAEA---E-LNKKVEALQNOVAEELEELSKLEDNL-KDAETNN-VEDYIK	366
BG7817	AAEA---E-LNKKVEALQNOVAEELEELSKLEDNL-KDAETNN-VEDYIK	366
Strain 18	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	122
BG9739	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AGOYL-	283
L81905	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AGOYL-	283
PspA AC94	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSGGGY-SALYL-	280
Strain 15	--E-----E-LSDKIDELDAETAKLEKDV---E-DF-KNSDGEQ-AEOYLV	145
ATCC6303	AAEDANIEALQNKVADLENKVAELDKEVTRLQSDL-KDAEENN-VEDYVK	348
BG7561	AAEA---E-LNKKVEALQNOVAEELEELSKLEDNL-KDAETNN-VEDYIK	366
Strain 23	AAEA---E-LDKKADELQNKVADLEKEISNLEILL-GGADPE---DDT--	133
Strain 18	AAEA---E-LDKKADELQNKVADLEKEISNLEILL-GGADPE---DDT--	135
SP 222	AAEDANIEALQNKVADLENKVAELDKEVTRLQSDL-KDAEENN-VEDYVK	127
BG9163	--E-----E-LSGKIEELDAETAELE--V---Q--L-KDAEGNNNVEAYFK	297
EF6796	--E-----E-LSGKIEELDAETAELE--V---Q--L-KDAEGNNNVEAYFK	296
BG8090	AAEA---E-LDKKADELQNKVADLEKEISNLEILL-GGADPE---DDT--	414
EF3296	AEFA---E-LDKKADELQNKVADLEKEISNLEILL-GGADSE---DDT--	413
Rx1	--E-----E-LSDKIDELDAETAKLE-D---Q--L-KAAEENNNVEDYFK	261
DBL1	--E-----E-LSDKIDELDAETAKLEKNV---EY-F-KKTDAEQ-TEOYL-	283
EF5668	AAEA---E-LNEKVEALQNOVAEELEELSKLEDNL-KDAETNN-VEDYIK	347
Wu2	--E-----E-LSDKIDELDAETAKLE-D---Q--L-KAVEENNNVEDYST	290
EF10197	--E-----E-LSDKIDELDAETAKLE-D---Q--L-KAAEENNNVEDYFK	282
PspA AC12	AEFA---E-LDKKADELQNKVADLEKEISNLEILL-GGADSE---DDT--	321
DBL5	--E-----E-LSGKIEELDAETAELE--V---Q--L-KDAEGNNNVEAYFK	284
DBL6A	--EEDRRNYPTNTYKTLEDAESDVKYKEAELELVKEEAKESRNEEKVK	366
BG6380	GEETLILA-IQNKISDLDDKIAEAEKKLADSQ-----NCEG-VEDYWT	303

Figure 6

Figure 7: CD Spectrum of CVX0270





Figure 8: Thermal Denaturation of Peptide CVX0270



## SEQUENCE LISTING

<110> Houston, Michael E.  
Hodges, Robert

<120> Use of Coiled-Coil Structural Scaffold to Generate  
Structure-Specific Peptides

<130> 003592-007

<150> US 60/211,892

<151> 2000-06-14

<150> US 60/213,387

<151> 2000-06-23

<160> 16

<170> PatentIn version 3.0

<210> 1

<211> 619

<212> PRT

<213> Streptococcus pneumoniae

<400> 1

Met	Asn	Lys	Lys	Lys	Met	Ile	Leu	Thr	Ser	Leu	Ala	Ser	Val	Ala	Ile
1				5					10					15	
Leu	Gly	Ala	Gly	Phe	Val	Ala	Ser	Gln	Pro	Thr	Val	Val	Arg	Ala	Glu
			20					25					30		
Glu	Ser	Pro	Val	Ala	Ser	Gln	Ser	Lys	Ala	Glu	Lys	Asp	Tyr	Asp	Ala
		35					40					45			
Ala	Lys	Lys	Asp	Ala	Lys	Asn	Ala	Lys	Lys	Ala	Val	Glu	Asp	Ala	Gln
	50					55					60				
Lys	Ala	Leu	Asp	Asp	Ala	Lys	Ala	Ala	Gln	Lys	Lys	Tyr	Asp	Glu	Asp
65					70					75				80	
Gln	Lys	Lys	Thr	Glu	Glu	Lys	Ala	Ala	Leu	Glu	Lys	Ala	Ala	Ser	Glu
			85						90					95	
Glu	Met	Asp	Lys	Ala	Val	Ala	Ala	Val	Gln	Gln	Ala	Tyr	Leu	Ala	Tyr
			100					105					110		



Gln	Gln	Ala	Thr	Asp	Lys	Ala	Ala	Lys	Asp	Ala	Ala	Asp	Lys	Met	Ile	
		115					120					125				
Asp	Glu	Ala	Lys	Lys	Arg	Glu	Glu	Glu	Ala	Lys	Thr	Lys	Phe	Asn	Thr	
	130					135					140					
Val	Arg	Ala	Met	Val	Val	Pro	Glu	Pro	Glu	Gln	Leu	Ala	Glu	Thr	Lys	
145					150					155					160	
Lys	Lys	Ser	Glu	Glu	Ala	Lys	Gln	Lys	Ala	Pro	Glu	Leu	Thr	Lys	Lys	
				165					170					175		
Leu	Glu	Glu	Ala	Lys	Ala	Lys	Leu	Glu	Glu	Ala	Glu	Lys	Lys	Ala	Thr	
			180					185					190			
Glu	Ala	Lys	Gln	Lys	Val	Asp	Ala	Glu	Glu	Val	Ala	Pro	Gln	Ala	Lys	
		195					200					205				
Ile	Ala	Glu	Leu	Glu	Asn	Gln	Val	His	Arg	Leu	Glu	Gln	Glu	Leu	Lys	
	210					215					220					
Glu	Ile	Asp	Glu	Ser	Glu	Ser	Glu	Asp	Tyr	Ala	Lys	Glu	Gly	Phe	Arg	
225					230					235					240	
Ala	Pro	Leu	Gln	Ser	Lys	Leu	Asp	Ala	Lys	Lys	Ala	Lys	Leu	Ser	Lys	
				245					250					255		
Leu	Glu	Glu	Leu	Ser	Asp	Lys	Ile	Asp	Glu	Leu	Asp	Ala	Glu	Ile	Ala	
			260					265					270			
Lys	Leu	Glu	Asp	Gln	Leu	Lys	Ala	Ala	Glu	Glu	Asn	Asn	Asn	Val	Glu	
		275					280					285				
Asp	Tyr	Phe	Lys	Glu	Gly	Leu	Glu	Lys	Thr	Ile	Ala	Ala	Lys	Lys	Ala	
	290					295					300					
Glu	Leu	Glu	Lys	Thr	Glu	Ala	Asp	Leu	Lys	Lys	Ala	Val	Asn	Glu	Pro	
305					310					315					320	
Glu	Lys	Pro	Ala	Pro	Ala	Pro	Glu	Thr	Pro	Ala	Pro	Glu	Ala	Pro	Ala	
				325					330					335		
Glu	Gln	Pro	Lys	Pro	Ala	Pro	Ala	Pro	Gln	Pro	Ala	Pro	Ala	Pro	Lys	
			340					345					350			
Pro	Glu	Lys	Pro	Ala	Glu	Gln	Pro	Lys	Pro	Glu	Lys	Thr	Asp	Asp	Gln	
		355					360					365				

Gln	Ala	Glu	Glu	Asp	Tyr	Ala	Arg	Arg	Ser	Glu	Glu	Glu	Tyr	Asn	Arg
	370					375					380				
Leu	Thr	Gln	Gln	Gln	Pro	Pro	Lys	Ala	Glu	Lys	Pro	Ala	Pro	Ala	Pro
385					390					395					400
Lys	Thr	Gly	Trp	Lys	Gln	Glu	Asn	Gly	Met	Trp	Tyr	Phe	Tyr	Asn	Thr
				405					410					415	
Asp	Gly	Ser	Met	Ala	Thr	Gly	Trp	Leu	Gln	Asn	Asn	Gly	Ser	Trp	Tyr
			420					425					430		
Tyr	Leu	Asn	Ser	Asn	Gly	Ala	Met	Ala	Thr	Gly	Trp	Leu	Gln	Tyr	Asn
		435					440					445			
Gly	Ser	Trp	Tyr	Tyr	Leu	Asn	Ala	Asn	Gly	Ala	Met	Ala	Thr	Gly	Trp
	450					455					460				
Ala	Lys	Val	Asn	Gly	Ser	Trp	Tyr	Tyr	Leu	Asn	Ala	Asn	Gly	Ala	Met
465					470					475					480
Ala	Thr	Gly	Trp	Leu	Gln	Tyr	Asn	Gly	Ser	Trp	Tyr	Tyr	Leu	Asn	Ala
				485					490					495	
Asn	Gly	Ala	Met	Ala	Thr	Gly	Trp	Ala	Lys	Val	Asn	Gly	Ser	Trp	Tyr
			500					505					510		
Tyr	Leu	Asn	Ala	Asn	Gly	Ala	Met	Ala	Thr	Gly	Trp	Leu	Gln	Tyr	Asn
		515					520					525			
Gly	Ser	Trp	Tyr	Tyr	Leu	Asn	Ala	Asn	Gly	Ala	Met	Ala	Thr	Gly	Trp
	530					535					540				
Ala	Lys	Val	Asn	Gly	Ser	Trp	Tyr	Tyr	Leu	Asn	Ala	Asn	Gly	Ala	Met
545					550					555					560
Ala	Thr	Gly	Trp	Val	Lys	Asp	Gly	Asp	Thr	Trp	Tyr	Tyr	Leu	Glu	Ala
				565					570					575	
Ser	Gly	Ala	Met	Lys	Ala	Ser	Gln	Trp	Phe	Lys	Val	Ser	Asp	Lys	Trp
			580					585					590		
Tyr	Tyr	Val	Asn	Gly	Leu	Gly	Ala	Leu	Ala	Val	Asn	Thr	Thr	Val	Asp
		595					600					605			
Gly	Tyr	Lys	Val	Asn	Ala	Asn	Gly	Glu	Trp	Val					
	610					615									

<210> 2  
 <211> 40  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> stabilizing strand

<220>  
 <221> MOD\_RES  
 <222> (2)  
 <223> Nle

<400> 2

Cys Xaa Gly Gly Gly Glu Ile Glu Ala Leu Lys Lys Glu Ile Glu Ala  
 1 5 10 15

Leu Lys Lys Glu Ile Glu Ala Leu Lys Lys Glu Ile Glu Ala Leu Lys  
 20 25 30

Lys Glu Ile Glu Ala Leu Lys Lys  
 35 40

<210> 3  
 <211> 38  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> hybrid sequence

<220>  
 <221> MOD\_RES  
 <222> (2)  
 <223> Nle

<400> 3

Cys Xaa Gly Ile Glu Glu Leu Glu Lys Lys Ile Thr Glu Leu Lys Gln  
 1 5 10 15

Lys Ile Asp Ala Leu Glu Asn Gln Ile His Arg Leu Glu Gln Glu Ile  
 20 25 30

Lys Glu Leu Asp Glu Ser

35

<210> 4  
 <211> 38  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> hybrid sequence

<220>  
 <221> MOD\_RES  
 <222> (2)  
 <223> Nle

<400> 4

Cys Xaa Gly Leu Glu Glu Ala Glu Lys Lys Ala Thr Glu Ala Lys Gln  
 1 5 10 15

Lys Val Asp Ala Leu Glu Asn Gln Val His Arg Leu Glu Gln Glu Leu  
 20 25 30

Lys Glu Ile Asp Glu Ser  
 35

<210> 5  
 <211> 20  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> consensus sequence

<220>  
 <221> PEPTIDE  
 <222> (4)..(20)  
 <223> Amino acids 4, 5, 12, 16 and 20 are Xaa wherein Xaa = any a  
 mino acid

<400> 5

Glu Glu Leu Xaa Xaa Lys Ile Asp Glu Leu Asp Xaa Glu Ile Ala Xaa  
 1 5 10 15

Leu Glu Lys Xaa  
 20

<210> 6  
 <211> 8  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> consensus sequence

<400> 6

Glu Glu Leu Ser Asp Lys Ile Asp  
 1 5

<210> 7  
 <211> 27  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> hybrid sequence

<220>  
 <221> MOD\_RES  
 <222> (2)  
 <223> Nle

<400> 7

Cys Xaa Gly Glu Ile Glu Ala Leu Lys Lys Lys Ile Glu Glu Leu Ser  
 1 5 10 15

Asp Lys Ile Asp Glu Leu Glu Lys Glu Ile Lys  
 20 25

<210> 8  
 <211> 21  
 <212> PRT  
 <213> Haemophilus influenzae

<400> 8

Ile Lys Lys Val Leu Glu Ile Gly Leu Asn Met Ser Gln Glu Ala Ser  
 1 5 10 15

Asn Leu Thr Ser Ala  
 20

<210> 9  
<211> 27  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> hybrid sequence

<400> 9

Glu Ala Glu Ile Lys Lys Leu Leu Glu Ile Ile Leu Asn Leu Ser Gln  
1 5 10 15

Glu Ile Ser Asn Leu Thr Ser Ala Leu Lys Gly  
20 25

<210> 10  
<211> 20  
<212> PRT  
<213> Streptococcus pneumoniae

<400> 10

Leu Glu Lys Asp Val Glu Asp Phe Lys Asn Ser Asp Gly Glu Gln Ala  
1 5 10 15

Glu Gln Val Leu  
20

<210> 11  
<211> 15  
<212> PRT  
<213> Streptococcus pneumoniae

<400> 11

Leu Glu Asp Asn Leu Lys Asp Ala Glu Thr Asn Asn Val Glu Asp  
1 5 10 15

<210> 12  
<211> 10  
<212> PRT  
<213> Streptococcus pneumoniae

<400> 12

Leu Ala Lys Lys Gln Thr Glu Leu Glu Lys  
1 5 10

<210> 13  
<211> 35  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> hybrid sequence

<400> 13

Asp Val Glu Asp Phe Lys Asn Ser Asp Gly Glu Leu Glu Asp Asn Leu  
1 5 10 15

Lys Asp Ala Glu Thr Asn Asn Val Glu Asp Ala Lys Lys Gln Thr Glu  
20 25 30

Leu Glu Lys  
35

<210> 14  
<211> 35  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> hybrid sequence

<400> 14

Asp Ile Glu Asp Leu Lys Asn Ser Ile Gly Glu Leu Glu Asp Asn Ile  
1 5 10 15

Lys Glu Leu Glu Thr Asn Ile Val Glu Leu Ala Lys Lys Ile Thr Glu  
20 25 30

Leu Glu Lys  
35

<210> 15  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Coiled-coil scaffold

<220>



<221> PEPTIDE  
<222> (1)  
<223> Amino acid 1 is Xaa wherein Xaa = A and A can be I, L or V.

<220>  
<221> PEPTIDE  
<222> (2)..(7)  
<223> Amino acids 2, 3, 5-7 are Xaa wherein Xaa = any amino acid.

<220>  
<221> PEPTIDE  
<222> (4)  
<223> Amino acid 4 is Xaa wherein Xaa = D and D can be I, L or V.

<220>  
<221> PEPTIDE  
<222> (1)..(7)  
<223> Amino acids 1-7 can be repeated one or more times.

<400> 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
1 5

<210> 16  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Coiled-coil scaffold

<220>  
<221> MOD\_RES  
<222> (2)  
<223> Nle

<220>  
<221> PEPTIDE  
<222> (5)..(10)  
<223> Amino acids 5, 6, and 8-10 are Xaa wherein Xaa = any amino acid.

<220>  
<221> PEPTIDE  
<222> (4)..(10)  
<223> Amino acids 4-10 can be repeated one or more times.

<400> 16

Cys	Xaa	Gly	Ile	Xaa	Xaa	Leu	Xaa	Xaa	Xaa
1				5					10